arXiv:1601.06692v2 [math.DS] 16 Aug 2016

ON TONELLI PERIODIC ORBITS WITH LOW ENERGY ON SURFACES

LUCA ASSELLE AND MARCO MAZZUCCHELLI

Abstract. We prove that, on a closed surface, a Lagrangian system defined by a Tonelli Lagrangian L possesses a periodic orbit that is a local minimum of the free-period action functional on every energy level belonging to the low range of energies $(e_0(L), c_0(L))$. We also prove that almost every energy level in $(e_0(L), c_{\rm u}(L))$ possesses infinitely many periodic orbits. These statements extend two results, respectively due to Taimanov and Abbondandolo-Macarini-Mazzucchelli-Paternain, known for the special case of electromagnetic Lagrangians.

Contents

1. Introduction 1
Acknowledgments 3
2. Local minimizers of the free-period action functional 4
2.1. Unique free-time local minimizers 4
2.2. Embedded global minimizers of the free-period action functional 7
2.3. Compactness of minimizing sequences 13
3. Free-period discretizations 25
3.1. Unique action minimizers 25
3.2. The discrete free-period action functional 27
3.3. The discrete least action principle 28
3.4. The Hessian of the free-period action functional 29
3.5. The iteration map 32
4. Multiplicity of low energy periodic orbits 36
4.1. A property of high iterates of periodic orbits 36
4.2. The minimax scheme 39
5. A Tonelli Lagrangian with few periodic orbits on low energy levels 41
6. Tonelli setting with a general magnetic form 42
References 44

1. Introduction

Hamiltonian systems on cotangent bundles sometimes admit a dual description as Lagrangian systems, which are second order dynamical systems on the configuration space. This is the case for geodesic flows, which can be viewed as Reeb flows on unit-sphere cotangent bundles, or equivalently as the family of curves γ on the configuration space satisfying the geodesic equation $\nabla_t\dot\gamma \equiv 0$. The widest class of fiberwise convex Hamiltonian functions admitting a dual Lagrangian is the

Date: January 25, 2016. *Revised*: August 16, 2016.
2000 *Mathematics Subject Classification.* 37J45, 58E05.
Key words and phrases. Tonelli Lagrangians, periodic orbits, Mañé critical values.

Tonelli one: a Hamiltonian $H : \mathrm{T}^*M \to \mathbb{R}$ is said to be Tonelli when its restriction to any fiber $H|_{\mathrm{T}^*_q M}$ is a superlinear function with everywhere positive-definite Hessian; the class of Lagrangians dual to the Tonelli Hamiltonians is given by the so-called Tonelli Lagrangians $L : \mathrm{T}M \to \mathbb{R}$, which satisfy the analogous properties as H on the tangent bundle. The name of these Lagrangians is due to the seminal work of Tonelli [Ton34], who established the existence and regularity of action minimizers joining a given pair of points in the configuration space. More modern applications of 1-dimensional calculus of variations, notably Aubry-Mather theory [Mat91a, Mat91b, Mat93] and weak KAM theory [Fat97], also concerns Lagrangians and Hamiltonians of Tonelli type. The background and further applications of Tonelli systems can be found in, e.g., [Fat08, CI99, Maz12, Sor15]. The purpose of this paper is to extend to the Tonelli class two important existence results for periodic orbits, which were proved for the smaller class of electromagnetic Lagrangians.

Given a closed manifold M and a Tonelli Lagrangian $L : \mathrm{T}M \to \mathbb{R}$, the associated Euler-Lagrange flow $\phi_L^t : \mathrm{T}M \to \mathrm{T}M$ is defined by $\phi_L^t(\gamma(0), \dot\gamma(0)) = (\gamma(t), \dot\gamma(t))$, where $\gamma : \mathbb{R} \to M$ is a smooth solution of the Euler-Lagrange equation, which in local coordinates can be written as

$$\tfrac{\mathrm{d}}{\mathrm{d}t} L_v(\gamma, \dot\gamma) - L_q(\gamma, \dot\gamma) = 0. \tag{1.1}$$

The energy function $E : \mathrm{T}M \to \mathbb{R}$, given by $E(q, v) = L_v(q, v)v - L(q, v)$, is a first integral for the Euler-Lagrange flow. The periodic orbits of the Euler-Lagrange flow contained in the energy hypersurface $E^{-1}(k)$ are in one to one correspondence with the critical points of the free-period action functional

$$\mathcal{S}_k : W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty) \to \mathbb{R},$$
$$\mathcal{S}_k(\Gamma, \tau) = \tau \int_0^1 \Big[L(\Gamma(t), \dot\Gamma(t)/\tau) + k \Big] \, \mathrm{d}t.$$

We see the pair (Γ, τ) as the τ-periodic curve $\gamma(t) := \Gamma(t/\tau)$, so that $\mathcal{S}_k(\Gamma, \tau)$ represents the action of γ at level k. In this paper, a crucial role will be played by the energy values $e_0(L) \le c_{\mathrm{u}}(L) \le c_0(L)$, which are given by

$$e_0(L) = \max_{q \in M} E(q, 0),$$
$$c_{\mathrm{u}}(L) = -\inf \left\{ \frac{1}{\tau} \int_0^\tau L(\gamma(t), \dot\gamma(t)) \, \mathrm{d}t \;\middle|\; \gamma : \mathbb{R}/\tau\mathbb{Z} \xrightarrow{C^\infty} M \text{ contractible} \right\},$$
$$c_0(L) = -\inf \left\{ \frac{1}{\tau} \int_0^\tau L(\gamma(t), \dot\gamma(t)) \, \mathrm{d}t \;\middle|\; \gamma : \mathbb{R}/\tau\mathbb{Z} \xrightarrow{C^\infty} M \text{ null-homologous} \right\}.$$

In the literature, $c_{\mathrm{u}}(L)$ and $c_0(L)$ are known as the Mañé critical values of the universal cover and of the abelian cover respectively. Alternative equivalent definitions of such values will be given in Sections 2.2 and 4.2 (more details can be found in [Con06, Abb13] and in references therein).

Our first result is the following.

Theorem 1.1. *Let M be a closed surface, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian. For every energy level $k \in (e_0(L), c_0(L))$, the Lagrangian system of L possesses a periodic orbit γ_k with energy k that is a local minimizer of the free-period action functional $\mathcal{S}_k$ over the space of absolutely continuous periodic curves. Moreover, $\mathcal{S}_k(\gamma_k) < 0$ and γ_k lifts to a simple closed curve in some finite covering of M.*

This result was first established by Taimanov [Tai92b, Tai91] for electromagnetic Lagrangians. His beautiful observation is that the free-period action functional, which is not bounded from below in the energy range concerned by the statement, becomes bounded from below when restricted to a suitable space of null-homologous, embedded, periodic multicurves. This latter space is not compact, but a clever argument of Taimanov shows that the global minima of the free period action functional are contained in the interior of the space, and as such are critical points. Unfortunately, the papers [Tai92b, Tai91] are rather short and some crucial details are only sketched. An alternative proof based on a regularity theorem for almost minimal currents in the sense of Almgren, a deep result from geometric measure theory, was provided later by Contreras, Macarini, and Paternain [CMP04]. Our proof of Theorem 1.1, which will be carried out in Section 2, fills the details in Taimanov's arguments, while at the same time generalizes the result to the class of Tonelli Lagrangians.

Our second result is the following.

Theorem 1.2. *Let M be a closed surface, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian. For almost every energy level k in the interval $(e_0(L), c_{\mathrm{u}}(L))$ the Lagrangian system of L possesses infinitely many periodic orbits with energy k and negative action.*

For the special case of electromagnetic Lagrangians, this result was proved by the second author together with Abbondandolo, Macarini, and Paternain in the recent paper [AMMP14] (see also [AMP15] for a previous preliminary result in this direction). The proof involves a highly non-trivial argument of local critical point theory for the free-period action functional. Such functional, in the classical $W^{1,2}$ functional setting, is C^∞ for the case of electromagnetic Lagrangians, but is not C^2 for general Tonelli Lagrangians. This poses major difficulties, as the local arguments from critical point theory involve in an essential way the Hessian of the functional. Our way to circumvent this lack of regularity is to develop a finite dimensional functional setting, which may have independent interest, inspired by Morse's broken geodesics approximation of path spaces [Mil63, Section 16]. The free-period action functional in such finite dimensional setting, which we will call the discrete free-period action functional, is C^∞. This will allow to carry out the proof of Theorem 1.2 along the line of [AMMP14]. The construction of the finite dimensional functional setting will be given in Section 3, and the proof of Theorem 1.2 in Section 4

Both Theorems 1.1 and 1.2 concern energy levels above $e_0(L)$. One may wonder whether the assertions of the theorems are still valid below $e_0(L)$. This is not case. Indeed, in Section 5 we will provide an example of Tonelli Lagrangian L such that $e_0(L) > \min E$ and, for all energy values $k > \min E$ sufficiently close to $\min E$, the corresponding Lagrangian system possesses only two periodic orbits with energy k, and none of such orbits is a local minimum of the free-period action functional.

Finally, Theorems 1.1 and 1.2 can be extended to the case where the Euler-Lagrange dynamics, or actually the dual Hamiltonian dynamics, is modified by a non-exact magnetic term. The precise setting and the statements will be provided in Section 6.

Acknowledgments. The authors are grateful to Gabriele Benedetti for pointing out an inaccuracy in the first draft. L.A. is partially supported by the DFG grant AB

360/2-1 "Periodic orbits of conservative systems below the Mañé critical energy value". M.M. is partially supported by the ANR projects WKBHJ (ANR-12-BS01-0020) and COSPIN (ANR-13-JS01-0008-01). Part of the work presented in this paper was carried out in November 2015 at the Ruhr-Universität Bochum. M.M. wishes to thank Alberto Abbondandolo for the kind hospitality, and the Mathematics Department in Bochum for providing a stimulating working environment.

2. Local minimizers of the free-period action functional

2.1. **Unique free-time local minimizers.** Let M be a closed manifold equipped with an arbitrary Riemannian metric g, which induces a distance $\mathrm{dist} : M \times M \to [0, \infty)$. Without loss of generality, we can assume that M is orientable: if this is not the case, we replace it by its orientation double covering. We consider a Tonelli Lagrangian $L : \mathrm{T}M \to \mathbb{R}$. We recall that such L is a smooth function whose restriction to any fiber of $\mathrm{T}M$ is superlinear with everywhere positive-definite Hessian. We are interested in the Euler-Lagrange dynamics, that is, in the solutions $\gamma : \mathbb{R} \to M$ of the Euler-Lagrange equation (1.1). This type of dynamics is conservative: the energy $E(q, v) = L_v(q, v)v - L(q, v)$ is constant along the lift $t \mapsto (\gamma(t), \dot\gamma(t))$ of a solution γ of the Euler-Lagrange equation.

For all $q_0, q_1 \in M$, we denote by $\mathrm{AC}(q_0, q_1)$ the space of absolutely continuous curves connecting q_0 with q_1, that is the space of all $\gamma : [0, \tau] \to M$ such that $\tau \in (0, \infty)$, $\gamma(0) = q_0$, and $\gamma(\tau) = q_1$. For any constant $k \in \mathbb{R}$, the action of one such curve γ with respect to the Lagrangian $L + k$ is given by

$$\mathcal{S}_k(\gamma) = \int_0^\tau L(\gamma(t), \dot\gamma(t))\, \mathrm{d}t + k\tau.$$

It is well known that the space $\mathrm{AC}(q_0, q_1)$ has the structure of a Banach manifold: it can be seen as the product of the interval $(0, \infty)$ with the space of absolutely continuous curves from the interval $[0, 1]$ to M joining q_0 and q_1. In this setting, the principle of least action guarantees that the extremal points of $\mathcal{S}_k : \mathrm{AC}(q_0, q_1) \to \mathbb{R}$ are precisely those γ that are solution of the Euler-Lagrange equation with energy $E(\gamma, \dot\gamma) \equiv k$. Following Mañé [Mañ97, CDI97] we define the **Mañé action potential**

$$\Phi_k : M \times M \to \mathbb{R} \cup \{-\infty\},$$
$$\Phi_k(q_0, q_1) := \inf \big\{ \mathcal{S}_k(\gamma) \;\big|\; \gamma \in \mathrm{AC}(q_0, q_1) \big\}.$$

The **Mañé critical value** $c(L)$ is defined as the infimum of the set of $k \in \mathbb{R}$ such that the function Φ_k is everywhere finite. It is easy to see that $c(L)$ is a finite value: it must be larger than or equal to

$$e_0(L) := \max E\big(\cdot, 0\big).$$

If $k \geq c(L)$, the action potential Φ_k is a Lipschitz function vanishing on the diagonal submanifold of $M \times M$. If $k > c(L)$, Φ_k is strictly positive outside the diagonal, and the infimum in its definition is actually a minimum: for any pair of points $q_0, q_1 \in M$ there exists a curve $\gamma \in \mathrm{AC}(q_0, q_1)$ satisfying $\Phi_k(q_0, q_1) = \mathcal{S}_k(\gamma)$; such a γ is necessarily a solution of the Euler-Lagrange equation with energy k.

It will be useful for us to consider also the Hamiltonian characterization of the Mañé critical value, which goes as follows. Let $H : \mathrm{T}^*M \to \mathbb{R}$ be the Tonelli

Hamiltonian dual to L, that is

$$H(q,p) = \max_{v \in \mathrm{T}_q M} \Big[pv - L(q,v) \Big]. \tag{2.1}$$

The critical value $c(L)$ can be defined as the infimum of k such that there exists a smooth function $u : M \to \mathbb{R}$ that satisfies the Hamilton-Jacobi inequality

$$H\big(q, \mathrm{d}u(q)\big) < k.$$

It is well known that, for any energy level $k > c(L)$, there exists a diffeomorphism of the cotangent bundle T^*M sending the energy hypersurface $H^{-1}(k)$ to the unit-sphere cotangent bundle of a Finsler metric F on M, see [CIPP98, Corollary 2]; in particular, the solutions of the Euler-Lagrange equation with energy k are reparametrizations of the geodesics of F, and the following holds.

Lemma 2.1. *Let $k > c(L)$. There exists $\tau_{\mathrm{inj}} > 0$ such that the following hold:*

(i) *Every point $q_0 \in M$ admits an open neighborhood $U_{q_0} \subset M$ such that the smooth map*

$$\psi_{q_0} : \big[0, \tau_{\mathrm{inj}}\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0} M\big) \to U_{q_0}$$

given by $\psi_{q_0}(\tau, v_0) = \pi \circ \phi_L^\tau(q_0, v_0)$ restricts to a diffeomorphism

$$\psi_{q_0} : \big(0, \tau_{\mathrm{inj}}\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0} M\big) \to U_{q_0} \setminus \{q_0\}.$$

(ii) *For each pair of points $q_0 \in M$ and $q_1 \in U_{q_0}$, let $(\tau, v_0) \in \psi_{q_0}^{-1}(q_1)$. The embedded curve $\gamma : [0, \tau] \hookrightarrow U_{q_0}$ given by $\gamma(t) := \psi_{q_0}(t, v_0)$ is the unique global minimizer of the action functional $\mathcal{S}_k : \mathrm{AC}(q_0, q_1) \to \mathbb{R}$.*

Proof. We already remarked that, since $k > c(L)$, there exists a Finsler metric F on M such that every solution of the Euler-Lagrange equation of L with energy k is a reparametrization of a unit-speed geodesic of F. Therefore, point (i) follows from the analogous statement about the exponential maps of closed Finsler manifolds (see, e.g., [AP94, BCS00]). As for point (ii), notice that it is enough to establish its assertion for all pair of points $q_0, q_1 \in M$ sufficiently close. The original assertion will follow by replacing τ_{inj} by a smaller positive constant.

For all $q_0, q_1 \in M$, we denote by $\gamma_{q_0,q_1} : [0, \tau_{q_0,q_1}] \to M$ a solution of the Euler-Lagrange equation with energy k such that $\mathcal{S}_k(\gamma_{q_0,q_1}) = \Phi_k(q_0, q_1)$. In particular, γ_{q_0,q_1} is a global minimizer of $\mathcal{S}_k : \mathrm{AC}(q_0, q_1) \to \mathbb{R}$. Notice that

$$\begin{aligned} \Phi_k(q_0,q_1) &= \mathcal{S}_{c(L)}(\gamma_{q_0,q_1}) + (k - c(L))\tau_{q_0,q_1} \\ &\geq \Phi_{c(L)}(q_0,q_1) + (k - c(L))\tau_{q_0,q_1}, \end{aligned}$$

and therefore

$$\tau_{q_0,q_1} \leq \Omega_k(q_0,q_1) := \frac{\Phi_k(q_0,q_1) - \Phi_{c(L)}(q_0,q_1)}{k - c(L)}. \tag{2.2}$$

The function $\Omega_k : M \times M \to [0, \infty)$ is Lipschitz and vanishes on the diagonal submanifold of $M \times M$. We denote by $\ell_k > 0$ the Lipschitz constant of Ω_k, so that

$$\Omega_k(q_0,q_1) = \underbrace{\Omega_k(q_0,q_0)}_{=0} + \underbrace{\Omega_k(q_0,q_1) - \Omega_k(q_0,q_0)}_{\leq \ell_k \operatorname{dist}(q_0,q_1)} \leq \ell_k \operatorname{dist}(q_0,q_1).$$

If $\operatorname{dist}(q_0, q_1) < \tau_{\mathrm{inj}}/\ell_k$, the estimate (2.2) implies that $\tau_{q_0,q_1} < \tau_{\mathrm{inj}}$, and therefore any global minimizer of the action functional $\mathcal{S}_k : \mathrm{AC}(q_0, q_1) \to \mathbb{R}$ is contained

in the open set U_{q_0}. Point (i) implies that such minimizer is unique and is an embedded curve. ∎

The following lemma shows that on any energy level in $(e_0(L), c(L)]$ the qualitative properties of the Euler-Lagrange dynamics are locally the same as on energy levels in $(c(L), \infty)$.

Lemma 2.2. *If $k > e_0(L)$, every point of the closed manifold M admits an open neighborhood W and a Tonelli Lagrangian $\tilde{L} : \mathrm{T}M \to \mathbb{R}$ such that $\tilde{L}|_{\mathrm{T}W} = L|_{\mathrm{T}W}$ and $c(\tilde{L}) < k$.*

Proof. Let $q_0 \in M$ be a given point. We choose a smooth function $u : M \to \mathbb{R}$ such that $\mathrm{d}u(q_0) = \partial_v L(q_0, 0)$. This implies that

$$H(q_0, \mathrm{d}u(q_0)) = -L(q_0, 0) \leq e_0(L) < k.$$

We choose an open neighborhood W of q_0 whose closure is contained in an open set $W' \subset M$ such that $H(q, \mathrm{d}u(q)) < k$ for all $q \in W'$. Let $\chi : M \to (0, 1]$ be a smooth bump function that is identically equal to 1 on W and is identically equal to a constant smaller than $k/\max\{H(q, \mathrm{d}u(q)) \,|\, q \in M\}$ outside W'. We define a new Tonelli Hamiltonian $\tilde{H} : \mathrm{T}^*M \to \mathbb{R}$ by $\tilde{H}(q, p) := \chi(q)\, H(q, p)$, and its dual Tonelli Lagrangian $\tilde{L} : \mathrm{T}M \to \mathbb{R}$. By construction, $\tilde{L}$ coincides with L on $\mathrm{T}W$, and we have $\tilde{H}(q, \mathrm{d}u(q)) < k$ for all $q \in M$. In particular, $k > c(\tilde{L})$. ∎

If $k < c(L)$, the action potential is identically equal to $-\infty$, and in particular there are no curves joining two given points that are global minimizers of the action. However, if $k > e_0(L)$ there are still local minimizers joining sufficiently close points.

Lemma 2.3. *Let $k > e_0(L)$. There exist $\tau_{\mathrm{inj}} > 0$ and $\rho_{\mathrm{inj}} > 0$ such that the following hold:*

(i) *Every point $q_0 \in M$ admits an open neighborhood $U_{q_0} \subset M$ containing the compact Riemannian ball $\overline{B_g(q_0, \rho_{\mathrm{inj}})} = \{q \in M \,|\, \mathrm{dist}(q_0, q) \leq \rho_{\mathrm{inj}}\}$ such that the smooth map*

$$\psi_{q_0} : \big[0, \tau_{\mathrm{inj}}\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0}M\big) \to U_{q_0}$$

given by $\psi_{q_0}(\tau, v_0) = \pi \circ \phi_L^\tau(q_0, v_0)$ restricts to a diffeomorphism

$$\psi_{q_0} : \big(0, \tau_{\mathrm{inj}}\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0}M\big) \to U_{q_0} \setminus \{q_0\}.$$

(ii) *For each pair of points $q_0 \in M$ and $q_1 \in U_{q_0}$, set $(\tau, v_0) := \psi_{q_0}^{-1}(q_1)$. The curve $\gamma : [0, \tau] \to U_{q_0}$, given by $\gamma(t) := \psi_{q_0}(t, v_0)$, is embedded and is the unique global minimizer of the restriction of the action functional $\mathcal{S}_k : \mathrm{AC}(q_0, q_1) \to \mathbb{R}$ to the open subset of absolutely continuous curves contained in U_{q_0}.*

In the following, we will say that the curve γ of point (ii) above is the **unique free-time local minimizer** with energy k joining q_0 and q_1.

Remark 2.4. If $k \leq e_0(L)$, the assertion of Lemma 2.2 is not valid for any point of M anymore, but only for those points $q_0 \in M$ such that $E(q_0, 0) < k$. Analogously, the assertions of Lemma 2.3 are still valid for all points q_0 such that $E(q_0, 0) < k$ and for some $\tau_{\mathrm{inj}} > 0$ depending on q_0. ∎

Proof of Lemma 2.3. By Lemma 2.2, we can find an open cover $W_1, ..., W_h$ of M and, for all $i = 1, ..., h$, a Tonelli Lagrangian $L_i : \mathrm{T}M \to \mathbb{R}$ such that $c(L_i) < k$ and $L_i|_{\mathrm{T}W_i} = L|_{\mathrm{T}W_i}$. Let $E_i : \mathrm{T}M \to \mathbb{R}$ be the associated energy functions

$$E_i(q, v) = \partial_v L_i(q, v)v - L_i(q, v).$$

We choose $r > 0$ large enough so that every energy hypersurface $E_i^{-1}(k)$ is contained in the ball tangent bundle of radius r, i.e.,

$$g_q(v, v) < r^2, \qquad \forall (q, v) \in E_i^{-1}(k).$$

We apply Lemma 2.1 to each Lagrangian L_i, thus obtaining the constant $\tau_i > 0$ and the maps

$$\psi_{i,q_0} : \big[0, \tau_i\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0}M\big) \to U_{i,q_0}.$$

We denote by $\rho_{\mathrm{leb}} > 0$ the Lebesgue number of the open cover $W_1, ..., W_h$. We recall that ρ_{leb} is such that, for every $q_0 \in M$, the open Riemannian ball $B_g(q_0, \rho_{\mathrm{leb}})$ is contained in some open set W_i of the cover. Notice that, for all $q_0 \in M$ and $i \in \{1, ..., h\}$, the open set U_{i,q_0} is contained in the Riemannian ball $B_g(q_0, \tau_i r)$. We reduce the positive constants τ_i, so that they all coincide to a same positive constant $\tau_{\mathrm{inj}} < \rho_{\mathrm{leb}}/r$. This implies that, for all $q_0 \in M$, there exists $i \in \{1, ..., h\}$ such that U_{i,q_0} is contained in the open set W_i, above which the Lagrangians L_i and L coincide; hence, we set $U_{q_0} := U_{i,q_0}$ and we define the map

$$\psi_{q_0} : \big[0, \tau_{\mathrm{inj}}\big) \times \big(E^{-1}(k) \cap \mathrm{T}_{q_0}M\big) \to U_{q_0},$$
$$\psi_{q_0}(\tau, v_0) := \pi \circ \phi_L^\tau(q_0, v_0) = \pi \circ \phi_{L_i}^\tau(q_0, v_0).$$

The claims of points (i-ii), except that $U_{q_0} \supset \overline{B_g(q_0, \rho_{\mathrm{inj}})}$, follow from Lemma 2.1 applied to the Lagrangian L_i. The inclusion $\overline{B_g(q_0, \rho_{\mathrm{inj}})} \subset U_{q_0}$ follows if we choose the constant ρ_{inj} such that

$$0 < \rho_{\mathrm{inj}} < \min\Big\{\mathrm{dist}\big(q_0, \pi \circ \phi_L^{\tau_{\mathrm{inj}}}(q_0, v_0)\big) \;\Big|\; (q_0, v_0) \in E^{-1}(k)\Big\}. \qquad \blacksquare$$

As a corollary of Lemma 2.3, we reobtain the following well known statement (see, e.g., [Abb13, Proof of Lemma 7.2] for the original proof).

Corollary 2.5. *Every absolutely continuous periodic curve $\gamma : \mathbb{R}/\tau\mathbb{Z} \to M$ with $\tau > 0$ and image contained in a Riemannian ball of diameter ρ_{inj} is contractible and satisfies $\mathcal{S}_k(\gamma) > 0$.*

Proof. We set $q_0 := \gamma(0)$. The curve γ is entirely contained in the Riemannian ball $B_g(q_0, \rho_{\mathrm{inj}})$, which in turn is contained in the open set U_{q_0} given by Lemma 2.3. Since U_{q_0} is contractible, γ is contractible as well. Consider the stationary curve $\gamma_0 : \{0\} \to \{q_0\}$. The curve γ might be stationary as well, but formally it is different from γ_0 since $\tau > 0$. By Lemma 2.3(ii), γ_0 is the unique global minimizer of the restriction of the action functional $\mathcal{S}_k : \mathrm{AC}(q_0, q_0) \to \mathbb{R}$ to the open subset of curves contained in U_{q_0}. Therefore $\mathcal{S}_k(\gamma) > \mathcal{S}_k(\gamma_0) = 0$. $\blacksquare$

2.2. **Embedded global minimizers of the free-period action functional.** From now on, we will assume that M is an orientable closed surface, that is,

$$\dim(M) = 2.$$

We will consider the functional $\mathcal{S}_k$ as defined on the space of absolutely continuous periodic curves on M with arbitrary period. The latter space can be identified with

the product $\mathrm{AC}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$, so that any pair (Γ, τ) in this product defines the τ-periodic curve $\gamma(t) := \Gamma(t/\tau)$. The functional $\mathcal{S}_k$ will be called the **free-period action**.

Let us consider the Mañé critical value $c_0(L)$. This can be defined in several equivalent ways: as the minimum of $c(L + \theta)$ where θ ranges among the closed 1-forms on M, as the Mañé critical value of the lift of L to the abelian covering of M, or as the minimum k such that the free-period action functional $\mathcal{S}_k$ is non-negative on the connected components of null-homologous periodic curves. It is easy to see that $e_0(L) \leq c_0(L)$. From now on, we fix an energy value

$$k \in (e_0(L), c_0(L)).$$

The proof of Theorem 1.1 will follow Taimanov's ideas [Tai91, Tai92b, Tai92a] for the electromagnetic case: the free-period action functional $\mathcal{S}_k$ is bounded from below on the space of embedded periodic multicurves that are the oriented boundary of a compact region of M; this suggests to look for the periodic orbit of Theorem 1.1 by minimizing $\mathcal{S}_k$ there. The major difficulty is that such a space of multicurves is not compact, and so might be the minimizing sequences.

Let us introduce a suitable space over which we will perform the minimization. For each positive integers m, we first define $\mathcal{C}(m)$ to be the space of multicurves $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m)$ such that

- (C1) for every $i \in \{1, ..., m\}$, the curve $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ is absolutely continuous and topologically embedded;
- (C2) any two curves γ_i and γ_j do not have mutual intersections;
- (C3) the multicurve $\boldsymbol{\gamma}$ is the oriented boundary of a possibly disconnected, embedded, oriented, compact surface $\Sigma \subset M$ whose orientation agrees with the one of M.

For all positive integers $m \leq n$, we then define $\mathcal{D}(m, n)$ to be the space of multicurves $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m)$ such that:

- (D1) each $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ is a continuous curve, and there exist
$$0 = \tau_{i,0} \leq \tau_{i,1} \leq ... \leq \tau_{i,n_i} = \tau_i$$
such that, for all $j = 0, ..., n_i - 1$,
$$\mathrm{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1})\big) \leq \rho_{\mathrm{inj}},$$
and the restriction $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ is the unique free-time local minimizer with energy k joining the endpoints (see the definition after Lemma 2.3);
- (D2) $n_1 + ... + n_m \leq n$.

We endow both $\mathcal{C}(m)$ and $\mathcal{D}(m, n)$ with the topologies induced by the inclusions into the space of absolutely continuous multicurves with m connected components (which is naturally diffeomorphic to the m-fold product of $\mathrm{AC}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$).

Lemma 2.6. *The space $\mathcal{D}(m, n)$ is compact.*

Proof. Let $\{\boldsymbol{\gamma}_\alpha = (\gamma_{\alpha,1}, ..., \gamma_{\alpha,m}) \mid \alpha \in \mathbb{N}\} \subseteq \mathcal{D}(m, n)$ be a sequence of multicurves, and $0 = \tau_{\alpha,i,0} \leq \tau_{\alpha,i,1} \leq ... \leq \tau_{\alpha,i,n_{\alpha,i}} = \tau_{\alpha,i}$ the time-decomposition for the connected component $\gamma_{\alpha,i}$ given in (D1). Since $n_{\alpha,1} + ... + n_{\alpha,m} \leq n$, by the pigeonhole principle we can extract a subsequence, which we still denote by $\{\boldsymbol{\gamma}_\alpha \mid \alpha \in \mathbb{N}\}$,

such that, for some $n_1, ..., n_m \in \mathbb{N}$, we have

$$n_{\alpha,i} = n_i, \qquad \forall \alpha \in \mathbb{N},\ i = 1, ..., m.$$

Since M is compact, up to extracting a further subsequence we can assume that, for each $i \in \{1, ..., m\}$ and $j \in \{1, ..., n_i\}$, the sequence $\{\gamma_{\alpha,i}(\tau_{\alpha,i,j}) \,|\, \alpha \in \mathbb{N}\}$ converges to some point $q_{i,j} \in M$. For each $i = 1, ..., m$ we set $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ to be the unique continuous curve such that, for some $0 = \tau_{i,0} \leq \tau_{i,1} \leq ... \leq \tau_{i,n_i} = \tau_i$, each portion $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ is the unique free-time local minimizer with energy k joining $q_{i,j}$ and $q_{i,j+1}$. The multicurve $\boldsymbol{\gamma} := (\gamma_1, ..., \gamma_m)$ clearly belongs to $\mathcal{D}(m, n)$, and $\boldsymbol{\gamma}_\alpha \to \boldsymbol{\gamma}$ as $\alpha \to \infty$. ∎

Finally, the space of multicurves that we will be interested in will be

$$\mathcal{M}(n) := \bigcup_{m=1}^{n} \mathcal{M}(m, n),$$

where

$$\mathcal{M}(m, n) := \mathcal{D}(m, n) \cap \overline{\mathcal{C}(m)}.$$

It readily follows from this definition and Lemma 2.6 that $\mathcal{M}(m, n)$ is compact, and so is $\mathcal{M}(n)$. With a slight abuse of notation, we will denote by $\mathcal{S}_k : \mathcal{M}(m, n) \to \mathbb{R}$ the free-period action functional with energy k on multicurves, given by

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \sum_{i=1}^{m} \mathcal{S}_k(\gamma_i) = \sum_{i=1}^{m} \left(\int_0^{\tau_i} L(\gamma_i(t), \dot{\gamma}_i(t))\, \mathrm{d}t + \tau_i k \right).$$

Clearly, $\mathcal{S}_k$ is continuous.

We have already seen in Corollary 2.5 that every absolutely continuous periodic curve $\gamma : \mathbb{R}/\tau\mathbb{Z} \to M$ contained in a Riemannian ball of diameter ρ_{inj} is contractible and satisfies $\mathcal{S}_k(\gamma) \geq 0$. On the other hand, up to passing to a finite covering of the configuration space M, we can always find embedded curves with negative action.

Lemma 2.7. *There exists a finite covering $M' \to M$ with the following property. If we lift the Tonelli Lagrangian L to $\mathrm{T}M'$, and denote by $\mathcal{M}'(m, n)$ and $\mathcal{S}'_k$ the associated spaces of multicurves and free-period action functional, then for n large enough there exists a periodic curve $\gamma \in \mathcal{M}'(1, n)$ with $\mathcal{S}'_k(\gamma) < 0$.*

Proof. Let M_0 be the abelian cover of our configuration space M, that is, the covering space whose fundamental group is the kernel of the Hurewicz homomorphism $\pi_1(M) \to H_1(M; \mathbb{Z})$. We lift the Tonelli Lagrangian L to a function $L_0 : \mathrm{T}M_0 \to \mathbb{R}$, and we denote by $\mathcal{M}^0(m, n)$ and $\mathcal{S}^0_k$ the associated spaces of multicurves and free-period action functional. Since $k < c_0(L)$, there exists an absolutely continuous loop $\gamma_0 : \mathbb{R}/\tau\mathbb{Z} \to M_0$ such that $\mathcal{S}^0_k(\gamma_0) < 0$. Let $n \in \mathbb{N}$ be large enough so that $\mathrm{dist}(\gamma_0(s), \gamma_0(t)) < \rho_{\mathrm{inj}}$ whenever $|s - t| \leq 1/n$, where $\rho_{\mathrm{inj}} > 0$ is the constant of Lemma 2.3(ii). We replace each portion $\gamma_0|_{[i/n, (i+1)/n]}$ of our curve with the unique free-time local minimizer with energy k joining the endpoints. The resulting curve, which we will still denote by $\gamma_0 : \mathbb{R}/\tau\mathbb{Z} \to M$ (possibly for a different period τ than before), is a piecewise solution of the Euler-Lagrange equation of L_0, and in particular is piecewise smooth. Up to perturbing the vertices $\gamma_0(i/n)$, for $i = 0, ..., k-1$, we can assume that γ_0 has only finitely many self-intersections, all of which are double points.

We claim that we can find such a γ_0 without self-intersections. Suppose that our γ_0 has at least one double point (otherwise we are done). Let $t_0 \in [0,\tau)$ be the smallest time such that $\gamma_0(t_0)$ is a double point of γ_0, and let $t_1 > t_0$ be the smallest time such that $\gamma_0(t_1) = \gamma_0(t_0)$. We define the piecewise smooth curves $\gamma_1 : \mathbb{R}/(t_1 - t_0)\mathbb{Z} \to M_0$ and $\gamma_2 : \mathbb{R}/(\tau - t_1 + t_0)\mathbb{Z} \to M_0$ by

$$\begin{aligned} \gamma_1(t) &= \gamma_0(t_0 + t), \qquad \forall t \in [0, t_1 - t_0], \\ \gamma_2(t) &= \gamma_0(t_1 + t), \qquad \forall t \in [0, \tau - t_1 + t_0]. \end{aligned}$$

Notice that

$$\begin{aligned} \gamma_0(\mathbb{R}/\tau\mathbb{Z}) &= \gamma_1(\mathbb{R}/(t_1 - t_0)\mathbb{Z}) \cup \gamma_2(\mathbb{R}/(\tau - t_1 + t_0)\mathbb{Z}), \\ \mathcal{S}^0_k(\gamma_0) &= \mathcal{S}^0_k(\gamma_1) + \mathcal{S}^0_k(\gamma_2), \end{aligned}$$

and that both γ_1 and γ_2 have strictly less number of double points than γ_0. Since $\mathcal{S}^0_k(\gamma_0) < 0$, we must have $\mathcal{S}^0_k(\gamma_1) < 0$ or $\mathcal{S}^0_k(\gamma_2) < 0$, say $\mathcal{S}^0_k(\gamma_1) < 0$. If γ_1 has no double points, we are done. Otherwise, we repeat the whole procedure with the curve γ_1. After a finite numbers of iterations of this procedure, we end up with a curve $\gamma' \in \mathcal{M}^0(1,k)$ such that $\mathcal{S}^0_k(\gamma') < 0$.

The group of Deck transformations of the abelian covering M_0 is, by definition, the first homology group $H_1(M;\mathbb{Z}) \simeq \mathbb{Z}^{b_1}$. Let $e_1, ..., e_{b_1}$ be the generators of $H_1(M;\mathbb{Z})$. For each vector $\boldsymbol{v} = (v_1, ..., v_{b_1}) \in \mathbb{Z}^{b_1}$, we consider the subgroup $H(\boldsymbol{v}) := \langle e_1^{v_1}, ..., e_{b_1}^{v_{b_1}} \rangle$ of $H_1(M)$. The quotient $H_1(M)/H(\boldsymbol{v})$ is a finite group, and therefore M admits a finite covering M' whose fundamental group is this quotient. The abelian covering M_0 is a covering of M'. If all the entries of the vector $\boldsymbol{v}$ are large enough, γ' is projected to a null-homologous curve γ in M' that is a piecewise solution of the Euler-Lagrange equation with energy k of the lift of L to $\mathrm{T}M'$, has no self-intersections (in particular $\gamma \in \mathcal{M}'(1,n)$), and satisfies $\mathcal{S}'_k(\gamma) < 0$. ∎

From now on, in order to simplify the notation, we will assume that $M = M'$, and we will denote by n_{neg} the minimal positive integer so that the conclusion of Lemma 2.7 holds for all $n \geq n_{\mathrm{neg}}$.

Let $n \geq n_{\mathrm{neg}}$. Since $\mathcal{M}(n)$ is compact and the free-period action functional $\mathcal{S}_k : \mathcal{M}(n) \to \mathbb{R}$ is continuous, there exists $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m) \in \mathcal{M}(n)$ such that

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \min_{\mathcal{M}(n)} \mathcal{S}_k < 0. \tag{2.3}$$

Notice that some connected component of $\boldsymbol{\gamma}$ may be a collapsed curve $\gamma_i : \{0\} \to M$. However, in this case $\mathcal{S}_k(\gamma_i) = 0$. Since $\mathcal{S}_k(\boldsymbol{\gamma}) < 0$, there exists a connected component γ_j of $\boldsymbol{\gamma}$ such that $\mathcal{S}_k(\gamma_j) < 0$. Such a γ_j is not a collapsed curve. Therefore, after removing all the collapsed connected components of $\boldsymbol{\gamma}$ we are left with a multicurve in $\mathcal{M}(n)$, which we still denote by $\boldsymbol{\gamma}$, that satisfies (2.3).

We say that $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ is **non-iterated** when τ_i is its minimal period.

Lemma 2.8. *Assume that $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m) \in \mathcal{M}(n)$ is a multicurve that satisfies (2.3) and has no connected component that is a collapsed curve. If in addition $\boldsymbol{\gamma} \in \mathcal{M}(n-1)$, then each of its connected components is a non-iterated periodic orbit of the Lagrangian system of L with energy k.*

Proof. Every connected component of $\boldsymbol{\gamma}$ has the form $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$, and there exists a sequence of time parameters $0 = \tau_{i,0} < \tau_{i,1} < ... < \tau_{i,n_i} = \tau_i$ such that



FIGURE 1. **(a)** The connected components γ_{i_h} intersecting at q. **(b)** The curve γ_i'.

each restriction $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]}$ is the unique free-time local minimizer with energy k joining the endpoints, which are distinct and lie at distance less than or equal to ρ_{inj}. Assume by contradiction that γ_{i_1} is not C^1 at some time τ_{i_1,j_1}. There may be other portions of the multicurve $\boldsymbol{\gamma}$ passing through $q := \gamma_{i_1}(\tau_{i_1,j_1})$. Thus, there are finitely many more (possibly zero) indices $i_2, ..., i_r \in \{1, ..., m\}$ and times $\tau_{i_2,j_2}, ..., \tau_{i_r,j_r}$ such that the (i_h, j_h)'s are pairwise distinct and satisfy

$$\gamma_{i_1}(\tau_{i_1,j_1}) = \gamma_{i_2}(\tau_{i_2,j_2}) = ... = \gamma_{i_r}(\tau_{i_r,j_r}) = q.$$

Up to replacing (i_1, j_1) with some of the other (i_h, j_h), we can assume that the curve γ_{i_1} is an "innermost" one around its corner $\gamma_{i_1}(\tau_{i_1,j_1})$. More precisely, this means that for an arbitrarily small open disk $B \subset M$ containing the point q the following holds. Let (a, b) be the widest interval containing τ_{i_1,j_1} and such that $\gamma_{i_1}|_{(a,b)}$ is contained in B. We denote by $B'' \subset B$ the connected component of $B \setminus \gamma_i|_{(a,b)}$ lying on the side of the obtuse angle formed by γ_i at q, and fix an arbitrary point q'' in the interior of B''. Consider a sequence of embedded multicurves $\boldsymbol{\gamma}_\alpha = (\gamma_{\alpha,1}, ..., \gamma_{\alpha,m}) \in \mathcal{C}(m)$ converging to $\boldsymbol{\gamma}$ as $\alpha \to \infty$. Let (a_α, b_α) be the widest interval containing τ_{i_1,j_1} and such that $\gamma_{\alpha,i_1}|_{(a_\alpha,b_\alpha)}$ is contained in B. We denote by B''_α the connected component of $B \setminus \gamma_{\alpha,i_1}|_{(a_\alpha,b_\alpha)}$ containing the point q''. Then, the "innermost" condition for γ_{i_1} is that, for all α large enough, all the points $\gamma_{\alpha,i_2}(\tau_{i_2,j_2}), ..., \gamma_{\alpha,i_r}(\tau_{i_r,j_r})$ belong to B''_α. The simple situation where the curves $\gamma_{i_1}, ..., \gamma_{i_r}$ have an isolated intersection at q is depicted in Figure 1(a).

We set $(i, j) := (i_1, j_1)$ and $B' := B \setminus B''$. Notice that a priori B' might have empty interior: this is the case when γ_i reaches the point q and goes back along the same path. Up to shrinking B around q, we can assume that the multicurve $\boldsymbol{\gamma}$ does not intersect the interior of B'. Let $\epsilon_1 > 0$ be small enough so that the restriction $\gamma_i|_{[\tau_{i,j}-\epsilon_1,\tau_{i,j}]}$ is contained in B' and has length less than the constant $\rho_{\text{inj}} > 0$ of Lemma 2.3. For $\epsilon_2 > 0$ sufficiently small, we remove the portion $\gamma_i|_{[\tau_{i,j}-\epsilon_1,\tau_{i,j}+\epsilon_2]}$ from γ_i and glue in the unique free-time local minimizer with energy k joining $\gamma_i(\tau_{i,j} - \epsilon_1)$ and $\gamma_i(\tau_{i,j} + \epsilon_2)$, see Figure 1(b). Lemma 2.3(i) guarantees that the portion that we glued in does not have self-intersections nor intersections with other points of γ_i except at the endpoints. We denote by γ_i' the modified curve, and by $\boldsymbol{\gamma}'$ the multicurve obtained from $\boldsymbol{\gamma}$ by replacing γ_i with γ_i'. Since $\boldsymbol{\gamma}$ belongs to

$\mathcal{M}(n-1)$ and since γ_i is an innermost curve around its corner $\tau_{i,j}$, $\boldsymbol{\gamma}'$ belongs to $\mathcal{M}(n)$. However, $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, since we replaced a non-smooth portion of γ_i with a local action minimizer with energy k. This contradicts (2.3), and shows that every connected component of $\boldsymbol{\gamma}$ is C^1, thus C^∞.

Finally, let us show that each periodic orbit $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ is non-iterated. Assume by contradiction that the minimal period of γ_i is τ_i' such that the quotient $p := \tau_i/\tau_i'$ is an integer larger than 1. Namely, γ_i is the p-th iterate of a periodic orbit. But on an orientable surface, every periodic curve that is sufficiently C^0-close to a p-th iterate must have a self-intersection. Consider again a sequence $\boldsymbol{\gamma}_\alpha = (\gamma_{\alpha,1}, ..., \gamma_{\alpha,m})$ in $\mathcal{C}(m)$ converging to $\boldsymbol{\gamma}$. For α large enough, the periodic curve $\gamma_{\alpha,i}$ has a self-intersection, which is impossible by the definition of $\mathcal{C}(m)$. ■

Lemma 2.9. *Assume that $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m) \in \mathcal{M}(n)$ is a multicurve that satisfies (2.3) and has no connected component that is a collapsed curve. If in addition $\boldsymbol{\gamma} \in \mathcal{M}(n-3)$, then $\boldsymbol{\gamma}$ is embedded in M.*

Proof. Lemma 2.8 tells us that the connected components of $\boldsymbol{\gamma}$ are non-iterated periodic orbits of L with energy k. In particular they are immersed curves in M, for the energy hypersurface $E^{-1}(k)$ does not intersect the zero-section of $\mathrm{T}M$. Moreover, the only (self or mutual) intersections among connected components of $\boldsymbol{\gamma}$ are tangencies. Assume by contradiction that there is at least a tangency q at some point along $\boldsymbol{\gamma}$, and consider all the branches of $\boldsymbol{\gamma}$ involved in this tangency: thus, there are only finitely many indices $i_1, ..., i_r \in \{1, ..., m\}$ and, for $h = 1, ..., r$, times $t_h \in \mathbb{R}/\tau_{i_h}\mathbb{Z}$ such that the (i_h, t_h)'s are pairwise distinct and satisfy

$$\gamma_{i_1}(t_1) = \gamma_{i_2}(t_2) = ... = \gamma_{i_r}(t_r) = q.$$

Consider a sequence of embedded multicurves $\boldsymbol{\gamma}_\alpha = (\gamma_{\alpha,1}, ..., \gamma_{\alpha,m}) \in \mathcal{C}(m)$ converging to $\boldsymbol{\gamma}$ as $\alpha \to \infty$. Up to permuting the indices $i_1, ..., i_r$ and extracting a subsequence of $\{\boldsymbol{\gamma}_\alpha \,|\, \alpha \in \mathbb{N}\}$ we can assume that the curves γ_{α,i_1} and γ_{α,i_2} are adjacent near the point q. More precisely, this means that there exists an arbitrarily small open disk $B \subset M$ such that, for all $\alpha \in \mathbb{N}$, the points $\gamma_{\alpha,i_1}(t_1)$ and $\gamma_{\alpha,i_2}(t_2)$ belong to the closure of a connected component B_α of $B \setminus \boldsymbol{\gamma}_\alpha$, see Figure 2(a).

The velocity vectors $\dot\gamma_{i_1}(t_1)$ and $\dot\gamma_{i_2}(t_2)$ are non-zero and parallel, i.e., $\dot\gamma_{i_1}(t_1) = \lambda\dot\gamma_{i_2}(t_2)$ for some real number $\lambda \neq 0$. We claim that $\lambda < 0$. Indeed, since the energy hypersurface $E^{-1}(k)$ intersects the fiber T_qM in a convex sphere, if $\lambda > 0$ then $\lambda = 1$, and therefore the periodic orbits γ_{i_1} and γ_{i_2} coincide. But then, for all α large enough, the curves γ_{α,i_1} and γ_{α,i_2} are the boundary of an open embedded annulus $A_\alpha \subset M$ containing the strip B_α and not containing any other connected component of $\boldsymbol{\gamma}_\alpha$. However, γ_{α,i_1} and γ_{α,i_2} are not the oriented boundary of A_α, since they are both oriented clockwise or counterclockwise. This contradicts the fact that the multicurve $\boldsymbol{\gamma}_\alpha$ is the oriented boundary of a compact embedded surface (condition (C3) in the definition of $\mathcal{C}(m)$). Now, consider the set

$$B' := \bigcap_{\beta\in\mathbb{N}} \overline{\bigcup_{\alpha>\beta} B_\alpha}.$$

The topological boundary $\partial B'$ is contained in $(\gamma_{i_1} \cup \gamma_{i_2}) \cap B$. Fix $\epsilon > 0$ small enough so that the curve $\gamma_{i_1}|_{[t_1-\epsilon, t_1]}$ has length less than ρ_{inj} and is contained in B'. For $\epsilon' > 0$ small enough, consider the unique action minimizer $\gamma' : [0, \tau'] \to M$ with energy k joining $\gamma'(0) = \gamma_{i_1}(t_1 - \epsilon)$ with $\gamma'(\tau') = \gamma_{i_2}(t_2 + \epsilon')$. Notice that γ' is



FIGURE 2. **(a)** The disk B and the strip B_α. **(b)** Outcome of the cut-and-paste procedure.

contained in $B' \setminus \{q\}$. We now modify the curves γ_{i_1} and γ_{i_2} as follows: along γ_{i_1}, once we reach $\gamma_{i_1}(t_1 - \epsilon)$, we continue along $\gamma'|_{[0,\tau']}$ and then along γ_{i_2}; along γ_{i_2}, once we reach $\gamma_{i_2}(t_2)$, we continue along γ_{i_1} (see Figure 2(b)).

If $i = i_1 = i_2$, this procedure replaces the connected component γ_i with two curves γ_i' and γ_i'' such that $\mathcal{S}_k(\gamma_i') + \mathcal{S}_k(\gamma_i'') < \mathcal{S}(\gamma_i)$; since $\boldsymbol{\gamma} \in \mathcal{M}(n-3)$, the multicurve $\boldsymbol{\gamma}'$ obtained from $\boldsymbol{\gamma}$ by replacing γ_i with $\gamma_i' \cup \gamma_i''$ belongs to $\mathcal{M}(n)$ and satisfies $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, which contradicts (2.3).

If $i_1 \neq i_2$, the procedure replaces the connected components γ_{i_1} and γ_{i_2} with a single curve $\gamma_{i_1 i_2}'$ such that $\mathcal{S}_k(\gamma_{i_1 i_2}') < \mathcal{S}_k(\gamma_{i_1}) + \mathcal{S}_k(\gamma_{i_2})$; since $\boldsymbol{\gamma} \in \mathcal{M}(n-3)$, the multicurve $\boldsymbol{\gamma}'$ obtained from $\boldsymbol{\gamma}$ by replacing $\gamma_{i_1} \cup \gamma_{i_2}$ with $\gamma_{i_1 i_2}'$ belongs to $\mathcal{M}(n)$ and satisfies $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, which contradicts (2.3). ∎

2.3. **Compactness of minimizing sequences.** The only missing ingredient to complete the proof of Theorem 1.1 is a compactness result: we wish to show that there exists $n_{\min} \in \mathbb{N}$ such that, for all $n \geq n_{\min}$, a minimizer of $\mathcal{S}_k$ over the space $\mathcal{M}(n)$ belongs to $\mathcal{M}(n_{\min})$, unless it contains collapsed connected components that we can always throw away. The proof of such statement will take most of this subsection. We begin with two preliminary lemmas.

Lemma 2.10. *Let $k > e_0(L)$. Consider piecewise smooth curves $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$, for $i = 1, ..., m$, such that the multicurve $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m)$ belongs to $\overline{\mathcal{C}(m)}$. Then*

$$\tau_1 + ... + \tau_m \leq \frac{\mathcal{S}_k(\boldsymbol{\gamma}) + \int_M |\mathrm{d}\theta|}{k - e_0(L)}. \tag{2.4}$$

Proof. Consider the smooth 1-form on M defined by $\theta_q(v) := L_v(q,0)v$, where L_v denotes the fiberwise derivative of L. The function $v \mapsto L(q,v) - \theta_q(v)$ has a global minimum at the origin, and therefore

$$L(q,v) - \theta_q(v) + e_0(L) \geq L(q,0) + e_0(L) = -E(q,0) + e_0(L) \geq 0.$$

We infer

$$\mathcal{S}_{e_0(L)}(\boldsymbol{\gamma}) - \int_{\boldsymbol{\gamma}} \theta = \sum_{i=0}^{m} \int_0^{\tau_i} \Big[L(\gamma_i, \dot{\gamma}_i) - \theta_{\gamma_i}(\dot{\gamma}_i) + e_0(L) \Big] \, \mathrm{d}t \geq 0. \tag{2.5}$$

Let $\{\boldsymbol{\gamma}_\alpha \mid \alpha \in \mathbb{N}\}$ be a sequence in $\mathcal{C}(m)$ converging to $\boldsymbol{\gamma}$ as $\alpha \to \infty$. If Σ_α is the compact submanifold whose oriented boundary is $\boldsymbol{\gamma}_\alpha$, we set

$$\Sigma := \bigcap_{\beta \in \mathbb{N}} \bigcup_{\alpha \geq \beta} \Sigma_\alpha.$$

We set $\tau := \tau_1 + ... + \tau_m$. By (2.5) and Stokes' Theorem, we have

$$\begin{aligned}
(k - e_0(L))\, \tau &\leq \mathcal{S}_{e_0(L)}(\boldsymbol{\gamma}) - \int_{\boldsymbol{\gamma}} \theta + (k - e_0(L))\, \tau \\
&= \mathcal{S}_k(\boldsymbol{\gamma}) - \int_{\boldsymbol{\gamma}} \theta \\
&= \mathcal{S}_k(\boldsymbol{\gamma}) - \int_{\Sigma} \mathrm{d}\theta \\
&\leq \mathcal{S}_k(\boldsymbol{\gamma}) - \int_M |\mathrm{d}\theta|,
\end{aligned}$$

which implies (2.4). ∎

Lemma 2.11. *Let $k > e_0(L)$. For all $s \in \mathbb{R}$ there exists $\ell_{\max}(s) > 0$ such that every $\boldsymbol{\gamma} \in \cup_{n \in \mathbb{N}} \mathcal{M}(n)$ satisfying $\mathcal{S}_k(\boldsymbol{\gamma}) \leq s$ has total length less than or equal to $\ell_{\max}(s)$.*

Proof. Since the energy hypersurface $E^{-1}(k)$ is compact, there exists $r > 0$ such that $g_q(v, v) \leq r^2$ for all $(q, v) \in E^{-1}(k)$. This, together with Lemma 2.10, implies that the length of $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m)$ can be bounded from above as

$$\begin{aligned}
\mathrm{length}(\boldsymbol{\gamma}) &= \sum_{i=0}^{m-1} \int_0^{\tau_i} \sqrt{g_{\gamma_i(t)}(\dot{\gamma}_i(t), \dot{\gamma}_i(t))} \, \mathrm{d}t \\
&\leq \big(\tau_1 + ... + \tau_m\big)\, r \\
&\leq \frac{\big(\mathcal{S}_k(\boldsymbol{\gamma}) + \int_M |\mathrm{d}\theta|\big)\, r}{k - e_0(L)} \\
&\leq \frac{\big(s + \int_M |\mathrm{d}\theta|\big)\, r}{k - e_0(L)} \\
&=: \ell_{\max}(s).
\end{aligned}$$

∎

In the following, we will be interested in the above lemma for $s = 0$, and we will simply write $\ell_{\max} := \ell_{\max}(0)$. We fix an integer $n \geq n_{\mathrm{neg}}$, and denote by $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m) \in \mathcal{M}(n)$ a multicurve such that

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \min_{\mathcal{M}(n)} \mathcal{S}_k. \tag{2.6}$$

By Lemma 2.7, up to replacing M by its finite covering M', we have that $\mathcal{S}_k(\boldsymbol{\gamma}) < 0$. As we did in the paragraph after equation (2.3), we remove from $\boldsymbol{\gamma}$ all the connected components that are collapsed curves, if there are any. This operation leaves the action $\mathcal{S}_k(\boldsymbol{\gamma})$ unchanged.

Lemma 2.12. *Each connected component of the multicurve $\boldsymbol{\gamma}$ has length larger than or equal to ρ_{inj}. In particular, there are at most $\ell_{\max}/\rho_{\text{inj}}$ many such connected components.*

Proof. Assume that some γ_i has length smaller than ρ_{inj}, and in particular it is contained in a Riemannian ball $B \subset M$ of diameter ρ_{inj}. Let $\gamma_{i_1}, \gamma_{i_2}, ..., \gamma_{i_r}$ be all the connected components of $\boldsymbol{\gamma}$ that are entirely contained in B. By Corollary 2.5, each γ_{i_j} is contractible and $\mathcal{S}_k(\gamma_{i_j}) > 0$. Therefore, the multicurve $\boldsymbol{\gamma}' := \boldsymbol{\gamma} \setminus \{\gamma_{i_1}, \gamma_{i_2}, ..., \gamma_{i_r}\}$ still belongs to $\mathcal{M}(n)$ and satisfies $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, which contradicts (2.6). This, together with the fact that the length of $\boldsymbol{\gamma}$ is at most $\ell_{\max}$, implies the lemma. ∎

Once $\boldsymbol{\gamma}$ is fixed, if needed, we reduce n so that

$$\boldsymbol{\gamma} \in \mathcal{M}(n) \setminus \mathcal{M}(n-1). \tag{2.7}$$

We write $\boldsymbol{\gamma} = (\gamma_1, ..., \gamma_m)$. We recall that, for each component $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$, there exist $0 = \tau_{i,0} < \tau_{i,1} < ... < \tau_{i,n_i} = \tau_i$ such that $\text{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1})\big) \leq \rho_{\text{inj}}$ and the restriction $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ is the unique free-time local minimizer with energy k joining the endpoints. Condition (2.7) implies $n_1 + ... + n_m = n$. We will call **vertices** the times $\tau_{i,j}$, and **segments** the portions $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$. The decomposition of the multicurve $\boldsymbol{\gamma}$ in segments is clearly not unique: for instance, if a connected component γ_i is smooth (and thus a periodic orbit of the Euler-Lagrange flow with energy k), we may be able to shift all the vertices around the curve. We say that a segment $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ is **short** if

$$\text{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1})\big) < \rho_{\text{inj}},$$

whereas we say that it is **long** if

$$\text{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1})\big) = \rho_{\text{inj}}.$$

We choose a decomposition of $\boldsymbol{\gamma}$ in segments so that:

- (S1) each smooth connected component γ_i contains at most one short segment;
- (S2) on each connected component γ_i that is not smooth, if $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ is a short segment then γ_i is not C^1 at $\tau_{i,j+1}$.

Lemma 2.13. *On each connected component γ_i that is not smooth, every short segment $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ contains a tangency of $\boldsymbol{\gamma}$, that is, there exists $h \in \{1, ..., m\}$ and $l \in \{0, ..., n_h\}$ such that $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ and $\gamma_h|_{[\tau_{h,l}, \tau_{h,l+1}]}$ are distinct segments of $\boldsymbol{\gamma}$ and have a mutual intersection.*

Proof. Assume by contradiction that the short segment $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ does not intersect any other segment of $\boldsymbol{\gamma}$. There exists an open neighborhood $B \subset M$ of $\gamma_i([\tau_{i,j}, \tau_{i,j+1}])$ such that no connected component of $\boldsymbol{\gamma}$ other than γ_i intersects B, and if $\gamma_i(t) \in B$ then $t \in (\tau_{i,j-1}, \tau_{i,j+2})$, see Figure 3(a). Since, by assumption, $\text{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1})\big) < \rho_{\text{inj}}$, for all $\epsilon > 0$ small enough we still have $\text{dist}\big(\gamma_i(\tau_{i,j}), \gamma_i(\tau_{i,j+1} + \epsilon)\big) < \rho_{\text{inj}}$. We denote by $\gamma'_\epsilon : [0, \omega_\epsilon] \to M$ the unique free-time local minimizer with energy k joining $\gamma'_\epsilon(0) = \gamma_i(\tau_{i,j})$ and $\gamma'_\epsilon(\omega_\epsilon) = \gamma_i(\tau_{i,j+1} + \epsilon)$. Up to further reducing ϵ such curve is entirely contained in B and, by Lemma 2.3, intersects $\gamma_i([\tau_{i,j}, \tau_{i,j+2}])$ only at the endpoints $\gamma'_\epsilon(0)$ and $\gamma'_\epsilon(\omega_\epsilon)$. However, γ'_ϵ may intersect $\gamma_i([\tau_{i,j-1}, \tau_{i,j}))$ (this happens for an arbitrarily small



FIGURE 3. **(a)** The neighborhood B of the segment $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]}$. **(b)** The modified curve γ'_i, with the portion $\gamma'_\epsilon|_{[\alpha_\epsilon,\omega_\epsilon]}$ glued in.

$\epsilon > 0$ only if γ_i has a cusp at $\tau_{i,j}$, that is, $\dot\gamma_i(\tau_{i,j}^-)$ and $\dot\gamma_i(\tau_{i,j}^+)$ are parallel and point in opposite directions). Consider the largest time $\alpha_\epsilon \in [0,\omega_\epsilon)$ such that $\gamma'_\epsilon(\alpha_\epsilon)$ lies on $\gamma_i((\tau_{i,j-1},\tau_{i,j}])$, and call $\sigma_\epsilon \in (\tau_{i,j-1},\tau_{i,j}]$ the unique time such that $q_0 := \gamma'_\epsilon(\alpha_\epsilon) = \gamma_i(\sigma_\epsilon)$. The restricted curve $\gamma'_\epsilon|_{[\alpha_\epsilon,\omega_\epsilon]}$ intersects $\boldsymbol\gamma$ only at the endpoints $\gamma'_\epsilon(\alpha_\epsilon),\gamma'_\epsilon(\omega_\epsilon)$ and, by Lemma 2.3, is still a unique free-time local minimizer with energy k. If ϵ is small enough, both curves $\gamma'_\epsilon|_{[\alpha_\epsilon,\omega_\epsilon]}$ and $\gamma_i|_{[\sigma_\epsilon,\tau_{i,j+1}+\epsilon]}$ are contained in the open set U_{q_0} given by Lemma 2.3 and join q_0 with $\gamma_i(\tau_{i,j+1}+\epsilon)$. Therefore

$$\int_{\alpha_\epsilon}^{\omega_\epsilon}\Big[L(\gamma'_\epsilon(t),\dot\gamma'_\epsilon(t))+k\Big]\,\mathrm{d}t < \int_{\sigma_\epsilon}^{\tau_{i,j+1}+\epsilon}\Big[L(\gamma_i(t),\dot\gamma_i(t))+k\Big]\,\mathrm{d}t. \tag{2.8}$$

We remove the portion $\gamma_i|_{[\sigma_\epsilon,\tau_{i,j+1}+\epsilon]}$ from γ_i and glue in the unique free-time local minimizer $\gamma'_\epsilon|_{[\alpha_\epsilon,\omega_\epsilon]}$. We denote by γ'_i the modified curve, and by $\boldsymbol\gamma'$ the multicurve obtained from $\boldsymbol\gamma$ by replacing γ_i with γ'_i, see Figure 3(b).

Clearly, $\boldsymbol\gamma'$ belongs to $\mathcal{M}(n)$. Furthermore, the inequality in (2.8) implies that $\mathcal{S}_k(\boldsymbol\gamma') < \mathcal{S}_k(\boldsymbol\gamma)$, which contradicts (2.6). ∎

We fix, once for all, a sequence $\{\boldsymbol\gamma_\alpha = (\gamma_{\alpha,1},...,\gamma_{\alpha,m}) \mid \alpha\in\mathbb{N}\} \subset \mathcal{C}(m)$ such that $\boldsymbol\gamma_\alpha \to \boldsymbol\gamma$ as $\alpha\to\infty$. Each multicurve $\boldsymbol\gamma_\alpha$ is without self-intersections, but a priori that is not the case for the limit curve $\boldsymbol\gamma$. Since more than two branches of $\boldsymbol\gamma$ may intersect (tangentially) at a same point, we need the following definition. We say that $(i_1,t_1),(i_2,t_2)$ is an **adjacent tangency** of $\boldsymbol\gamma$ at $q\in M$ when $(i_1,t_1)\neq(i_2,t_2)$, $q=\gamma_{i_1}(t_1)=\gamma_{i_2}(t_2)$, and for all sufficiently small neighborhoods $B\subset M$ of q and for all $\alpha\in\mathbb{N}$ large enough the points $\gamma_{\alpha,i_1}(t_1)$ and $\gamma_{\alpha,i_2}(t_2)$ belong to the closure of a same connected component of $B\setminus\boldsymbol\gamma_\alpha$. We will say that two segments $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]}$ and $\gamma_h|_{[\tau_{h,l},\tau_{h,l+1}]}$ contain a **mutual adjacent tangency**, and we will write

$$\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]} \asymp \gamma_h|_{[\tau_{h,l},\tau_{h,l+1}]},$$

if there exist $t_1\in[\tau_{i,j},\tau_{i,j+1}]$ and $t_2\in[\tau_{h,l},\tau_{h,l+1}]$ such that $(i,t_1),(h,t_2)$ is an adjacent tangency of $\boldsymbol\gamma$.

For all $i,h\in\{1,...,m\}$ we introduce the set

$$A_{i,h} := \big\{(j,l) \;\big|\; \gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]} \asymp \gamma_h|_{[\tau_{h,l},\tau_{h,l+1}]}\big\}.$$

We denote by $\#A_{i,h}$ the cardinality of this set.

Lemma 2.14. *The total number n of segments of the multicurve $\boldsymbol\gamma$ is bounded as*

$$n \le 2\frac{\ell_{\max}}{\rho_{\mathrm{inj}}} + 2\sum_{i\le h}\#A_{i,h}.$$

Proof. We denote by n_{long} and n_{short} the number of long segments and the number of short segments of $\boldsymbol{\gamma}$ respectively, so that $n = n_{\text{long}} + n_{\text{short}}$. The first one can be bounded as

$$n_{\text{long}} \leq \ell_{\max}/\rho_{\text{inj}}.$$

Let n'_{short} be the number of short segments in the smooth connected components of the multicurve $\boldsymbol{\gamma}$, and $n''_{\text{short}} := n_{\text{short}} - n'_{\text{short}}$ be the number of short segments in the non-smooth connected components. By our choice of the decomposition of $\boldsymbol{\gamma}$ in segments and by Lemma 2.12, we have

$$n'_{\text{short}} \leq m \leq \ell_{\max}/\rho_{\text{inj}}.$$

Finally, Lemma 2.13 implies that

$$n''_{\text{short}} \leq 2 \sum_{i \leq h} \#A_{i,h}. \qquad \blacksquare$$

In order to provide an upper bound for n, we are left to bound the cardinality of the set $A_{i,h}$. We first provide a uniform bound of the number of tangencies at an arbitrary given point $q \in M$. For $i = 1, ..., m$, we define

$$T_i(q) := \big\{ t \in \mathbb{R}/\tau_i\mathbb{Z} \;\big|\; \gamma_i(t) = q \big\}.$$

Lemma 2.15. *It $T_i(q)$ is not empty, let $t_0 < t_1 < ... < t_{u-1}$ be its ordered elements, with $0 \leq t_0 < t_{u-1} < \tau_i$. For each $j \in \mathbb{Z}/u\mathbb{Z}$, we see $[t_j, t_{j+1}]$ as a compact subset of the circle $\mathbb{R}/\tau_i\mathbb{Z}$, which is an interval if $u > 1$ or is the entire circle if $u = 1$. Each closed curve $\gamma_i|_{[t_j,t_{j+1}]}$ has length larger than or equal to ρ_{inj}. In particular,*

$$\#T_i(q) \leq \ell_{\max}/\rho_{\text{inj}}.$$

Proof. Assume by contradiction that, for some j, the curve $\gamma_i|_{[t_j,t_{j+1}]}$ has length smaller than ρ_{inj}. In particular, $\gamma_i([t_j, t_{j+1}]) \subseteq B(q, \rho_{\text{inj}}/2)$. By Lemma 2.3, for every $h = 1, ..., m$, if any connected component γ_h restricts to a loop $\gamma_h|_{[a,b]}$, for some $a < b$, that is contained in the Riemannian ball $B(q, \rho_{\text{inj}}/2)$ and satisfies $\gamma_h(a) = \gamma_h(b) = q$, then γ_h must possess a vertex $t \in [a, b]$; moreover, by Corollary 2.5, $\gamma_h|_{[a,b]}$ is contractible and satisfies $\mathcal{S}_k(\gamma_h|_{[a,b]}) > 0$; in this case, we cut the portion $\gamma_h|_{[a,b]}$ from γ_h and replace it by a vertex. We repeat this procedure iteratively as many times as possible, and we produce a multicurve $\boldsymbol{\gamma}'$ that still belongs to $\mathcal{M}(n)$ but satisfies $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, contradicting (2.6). This argument, together with the fact that γ_i has length at most $\ell_{\max}$, provides the desired upper bound for $\#T_i(q)$. $\blacksquare$

Now, for all pairs of distinct segments $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]} \asymp \gamma_h|_{[\tau_{h,l},\tau_{h,l+1}]}$, we fix a point $q_{(i,j),(h,l)} \in M$ at which the two segments have an adjacent tangency (if there are several such points, we choose one of them arbitrarily), thus forming the set

$$B_{i,h} := \big\{ q_{(i,j),(h,l)} \in M \;\big|\; (j,l) \in A_{i,h} \big\}.$$

Clearly, $\#B_{i,h} \leq \#A_{i,h}$. The following lemma states that also the opposite inequality holds, up to a multiplicative constant.

Lemma 2.16. $\#A_{i,h} \leq \#B_{i,h} \cdot (\ell_{\max}/\rho_{\text{inj}})^2$.

Proof. Consider the map $Q : A_{i,h} \to B_{i,h}$ given by $Q(j,l) = q_{(i,j),(h,l)}$, and an arbitrary $q \in B_{i,h}$. We denote by $\pi_1(j,l) = j$ and $\pi_2(j,l) = l$ the projections onto the first and second factors respectively. Notice that

$$\#(Q^{-1}(q)) \leq \#(\pi_1(Q^{-1}(q))) \cdot \#(\pi_2(Q^{-1}(q))).$$

The set $\pi_1(Q^{-1}(q))$ contains precisely all those $j \in \{1, ..., n_i\}$ such that the segment $\gamma_i|_{[\tau_{i,j}, \tau_{i,j+1}]}$ has an adjacent tangency at q with some segment of γ_h. Therefore, by Lemma 2.15, we have

$$\#(\pi_1(Q^{-1}(q))) \leq \#(T_i(q)) \leq \ell_{\max}/\rho_{\mathrm{inj}}.$$

Analogously

$$\#(\pi_2(Q^{-1}(q))) \leq \#(T_h(q)) \leq \ell_{\max}/\rho_{\mathrm{inj}}.$$

We conclude that

$$\begin{aligned} \#A_{i,h} &\leq \#B_{i,h} \cdot \max_{q \in B_{i,h}} \Big(\#(\pi_1(Q^{-1}(q))) \cdot \#(\pi_2(Q^{-1}(q))) \Big) \\ &\leq \#B_{i,h} \cdot (\ell_{\max}/\rho_{\mathrm{inj}})^2. \end{aligned}$$

■

Lemma 2.17. *For all distinct $i, h \in \{1, ..., m\}$, we have*

$$\#B_{i,h} \leq \max\Big\{ \big(\max\{2, \ell_{\max}/\rho_{\mathrm{inj}}\} + 1\big)!, \big(2\ell_{\max}/\rho_{\mathrm{inj}}\big)! \Big\}.$$

Proof. Let $0 \leq t_0 < ... < t_{u-1} < \tau_i$ and $s_0, ..., s_{u-1} \in [0, \tau_h)$ be such that:

- $B_{i,h} = \{\gamma_i(t_0), ..., \gamma_i(t_{u-1})\} = \{\gamma_h(s_0), ..., \gamma_h(s_{u-1})\}$;
- $\gamma_i(t_j) \neq \gamma_i(t_l)$ if $j \neq l$;
- $(i, t_j), (h, s_j)$ is an adjacent tangency of $\boldsymbol{\gamma}$ for each $j = 0, ..., u-1$.

At this point we employ a combinatorial statement due to Taimanov [Tai92b, Proposition 1]: for every finite set $W \subset \mathbb{R}$ with cardinality $\#W \geq w!$ and every injective map $f : W \to \mathbb{R}$ there exists a subset $W' \subset W$ of cardinality $\#W' \geq w$ such that $f|_{W'}$ is monotone. Let $w \in \mathbb{N}$ be the integer such that

$$w! \leq \#B_{i,h} < (w+1)!$$

The combinatorial statement implies that there exist $0 \leq j_0 < ... < j_{w-1} \leq \#B_{i,h}$ such that either

$$s_{j_0} < s_{j_1} < ... < s_{j_{w-1}} \tag{2.9}$$

or

$$s_{j_0} > s_{j_1} > ... > s_{j_{w-1}}. \tag{2.10}$$

Assume that (2.9) holds. For each $v \in \mathbb{Z}/w\mathbb{Z}$ we consider the closed curve

$$\zeta_v := \gamma_i|_{[t_{j_v}, t_{j_{v+1}}]} * \overline{\gamma_h|_{[s_{j_v}, s_{j_{v+1}}]}},$$

where $*$ denotes the concatenation of paths and the overline bar changes the orientation of a loop. We claim that, if $w > 2$, the loop ζ_v is not contractible. Indeed, assume that ζ_v is the boundary of a contractible compact subset $K \subset M$. Since γ_i and γ_h have adjacent tangencies at the points $q_v := \gamma_i(t_{j_v}) = \gamma_h(s_{j_v})$ and $q_{v+1} := \gamma_i(t_{j_{v+1}}) = \gamma_h(s_{j_{v+1}})$, the point $\gamma_h(s_{j_{v+2}})$ is forced to lie inside K. But then $\gamma_h(s_{j_{v+2}})$ must coincide with $\gamma_h(s_{j_v})$, as otherwise γ_i and γ_h could not have an adjacent tangency at the point $\gamma_i(t_{j_{v+2}}) = \gamma_h(s_{j_{v+2}})$, see Figure 4. Therefore



FIGURE 4. Situation under (2.9) with ζ_v contractible. The connected components γ_i and γ_h are the solid and dashed ones respectively. A priori, the compact set K may even have empty interior, but still the portion of γ_h that goes from q_{v+1} to q_v must be contained in K and cannot have an adjacent tangency with the portion of γ_i that goes from q_{v+1} to q_v.

$v = v + 2$ and $w = 2$. By Corollary 2.5, each loop ζ_v must have length larger than or equal to ρ_{inj}. Since

$$\sum_{v \in \mathbb{Z}/w\mathbb{Z}} \text{length}(\zeta_v) = \text{length}(\gamma_i) + \text{length}(\gamma_h) \leq \ell_{\max},$$

we conclude $w \leq \max\{2, \ell_{\max}/\rho_{\text{inj}}\}$, and therefore

$$\#B_{i,h} \leq \big(\max\{2, \ell_{\max}/\rho_{\text{inj}}\} + 1\big)!$$

Assume that (2.10) holds instead. For each $v \in \mathbb{Z}/w\mathbb{Z}$ we consider the closed curve

$$\xi_v := \gamma_i|_{[t_{j_v}, t_{j_{v+1}}]} * \gamma_h|_{[s_{j_v}, s_{j_{v+1}}]}.$$

We denote by w_{short} the number of closed curves ξ_v, for $v \in \mathbb{Z}/w\mathbb{Z}$, whose length is smaller than ρ_{inj}, and we set $w_{\text{long}} := w - w_{\text{short}}$. Since the total length of the multicurve $\boldsymbol{\gamma}$ is at most $\ell_{\max}$, we have $w_{\text{long}} \leq \ell_{\max}/\rho_{\text{inj}}$. We claim that

$$\#B_{i,h} \leq (2\ell_{\max}/\rho_{\text{inj}})!$$

Indeed, assume by contradiction that this inequality does not hold, so that $w > 2\ell_{\max}/\rho_{\text{inj}}$. This implies that $w_{\text{short}} > w_{\text{long}}$, and therefore there exists $v \in \mathbb{Z}/w\mathbb{Z}$ such that both ξ_v and ξ_{v+1} have length smaller than ρ_{inj}. We set, for $l = 0, 1, 2$,

$$q_{v+l} := \gamma_i(t_{j_{v+l}}) = \gamma_h(s_{j_{v+l}}) \in M.$$

By Corollary 2.5, the loops ξ_v and ξ_{v+1} are the boundary of some contractible compact subsets $K_v \subseteq B_g(q_v, \rho_{\text{inj}}/2)$ and $K_{v+1} \subseteq B_g(q_{v+1}, \rho_{\text{inj}}/2)$ respectively, and satisfy $\mathcal{S}_k(\xi_v) > 0$ and $\mathcal{S}_k(\xi_{v+1}) > 0$, see Figure 5(a). Corollary 2.5 further implies that no connected component of the multicurve $\boldsymbol{\gamma}$ is entirely contained in K_v or in K_{v+1}, as otherwise by removing from $\boldsymbol{\gamma}$ all these connected components we would obtain a new multicurve $\boldsymbol{\gamma}'$ still belonging to $\mathcal{M}(n)$ and satisfying $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\boldsymbol{\gamma})$, which would contradict (2.6). Notice that, by the very definition of the t_j's, there is at least one vertex of $\boldsymbol{\gamma}$ in $\gamma_i|_{[t_{j_v}, t_{j_{v+1}}]}$ or in $\gamma_h|_{[s_{j_{v+1}}, s_{j_v}]}$, and analogously there is at least one vertex of $\boldsymbol{\gamma}$ in $\gamma_i|_{[t_{j_{v+1}}, t_{j_{v+2}}]}$ or in $\gamma_h|_{[s_{j_{v+2}}, s_{j_{v+1}}]}$. We now modify the connected components γ_i and γ_j as follows: along γ_i, once we reach q_v, we continue along γ_h; along γ_h, once we reach q_{v+2}, we continue along γ_i; finally we remove the portions $\gamma_i|_{[t_{j_v}, t_{j_{v+2}}]}$ and $\gamma_h|_{[s_{j_{v+2}}, s_{j_v}]}$. This procedure replaces the connected components γ_i and γ_h with a single connected component $\gamma'_{i,h}$, see Figure 5(b). The multicurve $\boldsymbol{\gamma}' := (\boldsymbol{\gamma} \setminus \{\gamma_i, \gamma_h\}) \cup \gamma_{i,h}$ satisfies $\mathcal{S}_k(\boldsymbol{\gamma}') < \mathcal{S}_k(\gamma)$,



FIGURE 5. **(a)** The connected components γ_i and γ_h of the original multicurve $\boldsymbol{\gamma}$. **(b)** The connected component $\gamma'_{i,h}$ of the new multicurve $\boldsymbol{\gamma}'$.

and does not have more vertices than the original multicurve $\boldsymbol{\gamma}$. Moreover, since K_v and K_{v+1} does not contain entire connected components of $\boldsymbol{\gamma}$ and since each of the sets $\{(i,t_{j_v}),(h,s_{j_v})\}$, $\{(i,t_{j_{v+1}}),(h,s_{j_{v+1}})\}$, and $\{(i,t_{j_{v+2}}),(h,s_{j_{v+2}})\}$ is an adjacent tangency of $\boldsymbol{\gamma}$, the multicurve $\boldsymbol{\gamma}'$ belongs to $\overline{\mathcal{C}(m-1)}$, and therefore to $\mathcal{M}(n)$. This contradicts (2.6). ∎

Finding an upper bound for the cardinalities $\#B_{i,i}$ is a more difficult task, which requires some preliminaries. Given an absolutely continuous loop $\zeta : \mathbb{R}/\sigma\mathbb{Z} \to M$ such that $\zeta(t_0) = \zeta(t_1)$ for some $t_0, t_1 \in \mathbb{R}$ with $0 < t_1 - t_0 < \sigma$, we say that the **scission** of ζ at t_0, t_1 is the operation that produces the loops $\zeta' : \mathbb{R}/(t_1-t_0)\mathbb{Z} \to M$ and $\zeta'' : \mathbb{R}/(\sigma + t_0 - t_1)\mathbb{Z} \to M$ given by

$$\zeta'(t) = \zeta(t_0 + t), \qquad \forall t \in [0, t_1 - t_0],$$
$$\zeta''(t) = \zeta(t_1 + t), \qquad \forall t \in [0, \sigma + t_0 - t_1],$$

see Figure 6. Notice that $\mathcal{S}_k(\zeta) = \mathcal{S}_k(\zeta') + \mathcal{S}_k(\zeta'')$.

We say that a periodic curve $\zeta_0 : \mathbb{R}/\sigma\mathbb{Z} \to M$, with $\sigma > 0$, is a **tree of short loops** when, for increasing values of the integer j going from 1 to some $h \in \mathbb{N}$, a scission of ζ_{j-1} produces the loops ω_j and ζ_j, and each of the loops $\omega_1, \omega_2, ..., \omega_h, \omega_{h+1} := \zeta_h$ has length less than ρ_{inj}. In particular, each of these last $h+1$ loops is contained in a Riemannian ball of diameter ρ_{inj}. Therefore, by Corollary 2.5, we have

$$\mathcal{S}_k(\zeta_0) = \mathcal{S}_k(\omega_1) + \mathcal{S}_k(\omega_2) + ... + \mathcal{S}_k(\omega_h) + \mathcal{S}_k(\omega_{h+1}) > 0.$$

Given a multicurve $\boldsymbol{\gamma}' = (\gamma'_1, ..., \gamma'_{m'}) \in \mathcal{M}(n')$ with connected components of the form $\gamma'_i : \mathbb{R}/\tau'_i\mathbb{Z} \to M$, a finite subset $B' \subset M$, and a pair $t_0, t_1 \in \mathbb{R}$ with $0 < t_1 - t_0 < \tau'_i$, we say that the restriction $\gamma'_i|_{[t_0,t_1]}$ is B'**-minimal** if $(i,t_0),(i,t_1)$ is an adjacent tangency of $\boldsymbol{\gamma}'$ at some point of B', and for all $t' \in (t_0,t_1)$ and $t'' \in [t_0,t_1] \setminus \{t'\}$ such that $q := \gamma'_i(t') = \gamma'_i(t'')$ we have $q \notin B'$.

Now, consider a connected component $\gamma_i : \mathbb{R}/\tau_i\mathbb{Z} \to M$ of our multicurve $\boldsymbol{\gamma}$. We assume that $B_{i,i} \neq \varnothing$ (otherwise we are done). We wish to apply an iterative procedure to γ_i that decomposes it into $B_{i,i}$-minimal loops and threes of short loops based at points of $B_{i,i}$. Therefore, we set

$$\mu_0 := \gamma_i.$$

By Lemma 2.15, no scission of μ_0 at times t'_0, t'_1 such that $\mu_0(t'_0) = \mu_0(t'_1) \in B_{i,i}$ produces two loops one of which has length smaller than ρ_{inj}. Therefore, we can find a restriction $\mu_0|_{[t_0,t_1]}$ that is $B_{i,i}$-minimal and has length larger than or equal to ρ_{inj}. We perform a scission of μ_0 at t_0, t_1, thus obtaining two loops ζ_1 and ζ'_1, the first



FIGURE 6. The curve ζ, and the result of its scission at t_0, t_1.

one corresponding to $\mu_0|_{[t_0,t_1]}$. Notice that the multicurve $(\boldsymbol{\gamma} \setminus \mu_0) \cup \zeta_1 \cup \zeta_1'$ belongs to $\mathcal{M}(n+2)$, since the scission of a loop adds at most one vertex to each of the new connected components produced. Moreover $\mathcal{S}_k\big((\boldsymbol{\gamma} \setminus \mu_0) \cup \zeta_1 \cup \zeta_1'\big) = \mathcal{S}_k(\boldsymbol{\gamma})$. The time 0 is the vertex of the loop ζ_1' corresponding to the scission point $\zeta_1'(0) = \zeta_1(0) \in B_{i,i}$. We have two possible cases.

- If no restriction of ζ_1' to some interval $[s_0, s_1]$ is a tree of short loops such that $\zeta'(s_0) = \zeta'(s_1) \in B_{i,i}$, we set $\mu_1 := \zeta_1'$.
- Otherwise, we consider such an interval $[s_0, s_1]$, and notice that $0 \in [s_0, s_1]$. We require $[s_0, s_1]$ to be maximal, meaning that $[s_0, s_1]$ is not strictly contained in a larger interval $[r_0, r_1]$ such that $\zeta_1'|_{[r_0,r_1]}$ is a tree of short loops with $\zeta_1'(r_0) = \zeta_1'(r_1) \in B_{i,i}$. Notice that ζ_1' has an adjacent self-tangency at times s_0, s_1. We perform a scission of ζ_1' at times s_0, s_1, which produces a tree of short loops ω_1 and a loop μ_1.

In both cases, no scission of μ_1 at times t_0', t_1' such that $\mu_1(t_0') = \mu_1(t_1') \in B_{i,i}$ can produce a tree of short loops. We then repeat the whole procedure with the loop μ_1 (instead of μ_0). The algorithm will eventually terminate after $a \in \mathbb{N}$ iterations, and the outcome will be the loops $\zeta_1, ..., \zeta_a, \omega_1, ..., \omega_b$, for some $0 \leq b \leq a$.

Since every $B_{i,i}$-minimal loop ζ_j produced has length at least ρ_{inj}, we have $a \leq \ell_{\max}/\rho_{\text{inj}}$. Moreover, the multicurve

$$\boldsymbol{\zeta} := (\boldsymbol{\gamma} \setminus \gamma_i) \cup \zeta_1 \cup ... \cup \zeta_a \cup \omega_1 \cup ... \cup \omega_b$$

belongs to $\mathcal{M}(n + 2a + 2b) \subset \mathcal{M}(n + 4\ell_{\max}/\rho_{\text{inj}})$ and satisfies $\mathcal{S}_k(\boldsymbol{\zeta}) = \mathcal{S}_k(\boldsymbol{\gamma})$. In order to simplify the notation, we write

$$\boldsymbol{\zeta} = (\zeta_1, ..., \zeta_a, \zeta_{a+1}, ..., \zeta_{a+b}, \zeta_{a+b+1}, ..., \zeta_{a+b+n-1}),$$

where

$$(\zeta_{a+1}, ..., \zeta_{a+b}) := (\omega_1, ..., \omega_b),$$
$$(\zeta_{a+b+1}, ..., \zeta_{a+b+n-1}) := (\gamma_1, ..., \gamma_{i-1}, \gamma_{i+1}, ..., \gamma_n).$$

For each $h = 1, ..., a+b$, we consider a decomposition in segments of the connected component $\zeta_h : \mathbb{R}/\sigma_h\mathbb{Z} \to M$ that satisfies the conditions as in (S1-S2): we introduce the time-decomposition $0 = \sigma_{h,0} \leq \sigma_{h,1} \leq ... \leq \sigma_{h,m_h} = \sigma_h$, so that the restrictions $\zeta_h|_{[\sigma_{h,l},\sigma_{h,l+1}]}$ are the segments of ζ_h. For all pairs of distinct segments $\zeta_h|_{[\sigma_{h,l},\sigma_{h,l+1}]} \asymp \zeta_x|_{[\sigma_{x,y},\sigma_{x,y+1}]}$ we fix a point $p_{(h,l),(x,y)} \in M$ at which the two segments have an adjacent tangency; if possible, we fix such a point so that it belongs to $B_{i,i}$. We define the finite set

$$C_{h,x} := \left\{ p_{(h,l),(x,y)} \;\middle|\; 1 \leq l \leq m_h,\, 1 \leq y \leq m_x,\, \zeta_h|_{[\sigma_{h,l},\sigma_{h,l+1}]} \asymp \zeta_x|_{[\sigma_{x,y},\sigma_{x,y+1}]} \right\}$$

Notice that

$$B_{i,i} \subset \bigcup_{1\leq h\leq x\leq a+b} C_{h,x}. \tag{2.11}$$

Furthermore, since $\zeta_1, ..., \zeta_a$ are $B_{i,i}$-minimal loops, we have

$$B_{i,i} \cap \big(C_{1,1} \cup ... \cup C_{a,a}\big) = \varnothing. \tag{2.12}$$

The estimates for the cardinalities of the remaining sets $C_{h,x}$ are provided by the following two lemmas.

Lemma 2.18. *For all $h = 1, ..., b$ we have $\#C_{a+h,a+h} < (4\ell_{\max}/\rho_{\mathrm{inj}})^2$.*

Proof. We recall that we have denoted by m_{a+h} the number of segments of ζ_{a+h}, and therefore $\#C_{a+h,a+h} \leq m_{a+h}^2$. Assume by contradiction that $\#C_{a+h,a+h} \geq (4\ell_{\max}/\rho_{\mathrm{inj}})^2$. In particular, ζ_{a+h} contains at least $4\ell_{\max}/\rho_{\mathrm{inj}}$ segments. Since ζ_{a+h} is a tree of short loops, a sequence of scissions decomposes it into finitely many loops $\eta_1, ..., \eta_s$, each one having length smaller than ρ_{inj}. In particular, each η_j is the topological boundary of a contractible compact set $K_j \subset B_g(\eta_j(0), \rho_{\mathrm{inj}}/2)$. We remove from the multicurve $\boldsymbol{\zeta}$ all the connected components that are entirely contained in one of the K_j's, and we further remove the connected component ζ_{a+h}. We denote the obtained multicurve by $\boldsymbol{\zeta}'$. Since we have removed at least $4\ell_{\max}/\rho_{\mathrm{inj}}$ segments, $\boldsymbol{\zeta}'$ belongs to $\mathcal{M}(n)$ and satisfies

$$\mathcal{S}_k(\boldsymbol{\zeta}') \leq \mathcal{S}_k(\boldsymbol{\zeta}) - \mathcal{S}_k(\zeta_{a+h}) < \mathcal{S}_k(\boldsymbol{\zeta}) = \mathcal{S}_k(\boldsymbol{\gamma}),$$

which contradicts (2.6). ∎

Lemma 2.19. *For all distinct $h, x \in \{1, ..., a+b\}$, we have*

$$\#C_{h,x} \leq \max\Big\{\big(\max\{2, \ell_{\max}/\rho_{\mathrm{inj}}\}+1\big)!, \big(4(\ell_{\max}/\rho_{\mathrm{inj}})^2 + (\ell_{\max}/\rho_{\mathrm{inj}})\big)!\Big\} =: c.$$

Proof. The proof is analogous to the one of Lemma 2.17, but we provide full details for the reader's convenience. Let $0 \leq t_0 < ... < t_{u-1} < \sigma_h$ and $s_0, ..., s_{u-1} \in [0, \sigma_x)$ be such that:

- $C_{h,x} = \big\{\zeta_h(t_0), ..., \zeta_h(t_{u-1})\big\} = \big\{\zeta_x(s_0), ..., \zeta_x(s_{u-1})\big\}$;
- $\zeta_h(t_l) \neq \zeta_h(t_y)$ if $l \neq y$;
- $(h, t_j), (x, s_j)$ is an adjacent tangency of $\boldsymbol{\zeta}$ for each $j = 0, ..., u-1$.

Let $w \in \mathbb{N}$ be the integer such that $w! \leq \#C_{h,x} < (w+1)!$. The combinatorial statement [Tai92b, Proposition 1] implies that there exist $0 \leq j_0 < ... < j_{w-1} \leq \#C_{h,x}$ such that either

$$s_{j_0} < s_{j_1} < ... < s_{j_{w-1}} \tag{2.13}$$

or

$$s_{j_0} > s_{j_1} > ... > s_{j_{w-1}}. \tag{2.14}$$

Assume that (2.13) holds. For each $v \in \mathbb{Z}/w\mathbb{Z}$ we consider the closed curve

$$\xi_v := \zeta_h|_{[t_{j_v}, t_{j_{v+1}}]} * \overline{\zeta_x|_{[s_{j_v}, s_{j_{v+1}}]}}.$$

We claim that, if $w > 2$, the loop ξ_v is not contractible. Indeed, assume that ξ_v is the boundary of a contractible compact subset $K \subset M$. Since ζ_h and ζ_x have adjacent tangencies at the points $q_v := \zeta_h(t_{j_v}) = \zeta_x(s_{j_v})$ and $q_{v+1} := \zeta_h(t_{j_{v+1}}) = \zeta_x(s_{j_{v+1}})$,

the point $\zeta_x(s_{j_{v+2}})$ is forced to lie inside K. But then $\zeta_x(s_{j_{v+2}})$ must coincide with $\zeta_x(s_{j_v})$, as otherwise ζ_h and ζ_x could not have an adjacent tangency at the point $\zeta_h(t_{j_{v+2}}) = \zeta_x(s_{j_{v+2}})$. Therefore $v = v + 2$ and $w = 2$. By Corollary 2.5, each loop ξ_v must have length larger than or equal to ρ_{inj}. Since

$$\sum_{v\in\mathbb{Z}/w\mathbb{Z}} \text{length}(\xi_v) = \text{length}(\gamma_h) + \text{length}(\gamma_x) \leq \ell_{\max},$$

we conclude $w \leq \max\{2, \ell_{\max}/\rho_{\text{inj}}\}$, and therefore

$$\#C_{h,x} \leq \big(\max\{2, \ell_{\max}/\rho_{\text{inj}}\} + 1\big)!$$

Assume that (2.14) holds instead. For each $v \in \mathbb{Z}/w\mathbb{Z}$ we consider the closed curve

$$\xi_v := \zeta_h|_{[t_{j_v}, t_{j_{v+1}}]} * \zeta_x|_{[s_{j_v}, s_{j_{v+1}}]}.$$

We denote by w_{short} the number of closed curves ξ_v, for $v \in \mathbb{Z}/w\mathbb{Z}$, whose length is smaller than ρ_{inj}, and we set $w_{\text{long}} := w - w_{\text{short}}$. Since the total length of the multicurve $\boldsymbol{\zeta}$ is at most $\ell_{\max}$, we have $w_{\text{long}} \leq \ell_{\max}/\rho_{\text{inj}}$. We claim that

$$\#C_{h,x} \leq \big(4(\ell_{\max}/\rho_{\text{inj}})^2 + (\ell_{\max}/\rho_{\text{inj}})\big)!$$

Indeed, assume by contradiction that the above inequality does not hold. This implies

$$w_{\text{short}} > 4\left(\frac{\ell_{\max}}{\rho_{\text{inj}}}\right)^2 + \frac{\ell_{\max}}{\rho_{\text{inj}}} - w_{\text{long}} \geq 4\left(\frac{\ell_{\max}}{\rho_{\text{inj}}}\right)^2 \geq 4\left(\frac{\ell_{\max}}{\rho_{\text{inj}}}\right) w_{\text{long}}.$$

Therefore, there exists $v \in \mathbb{Z}/w\mathbb{Z}$ such that the loops $\xi_v, \xi_{v+1}, ..., \xi_z$, for $z := v - 1 + 4\ell_{\max}/\rho_{\text{inj}}$, are distinct and each one has length smaller than ρ_{inj}. By Corollary 2.5, each of these loops ξ_y is the boundary of a contractible compact subset $K_y \subset B_g(\xi_y(0), \rho_{\text{inj}}/2)$ and satisfies $\mathcal{S}_k(\xi_y) > 0$. Notice that, by the very definition of the t_j's, each loop ξ_y contains at least one vertex of $\boldsymbol{\zeta}$. We now modify the connected components ζ_h and ζ_x as follows: along ζ_h, once we reach $\zeta_h(t_{j_v})$, we continue along ζ_x; along ζ_x, once we reach $\zeta_x(s_{j_z})$, we continue along ζ_h; finally we throw away the portions $\zeta_h|_{[t_{j_v}, t_{j_z}]}$ and $\zeta_x|_{[s_{j_z}, s_{j_v}]}$. This procedure replaces ζ_h and ζ_x with a single connected component $\zeta'_{h,x}$. We define the multicurve

$$\boldsymbol{\zeta}' := \big(\boldsymbol{\zeta} \setminus (\zeta_h \cup \zeta_x)\big) \cup \zeta'_{h,x}.$$

We remove from $\boldsymbol{\zeta}'$ all the connected components that are entirely contained in a compact set among $K_v, K_{v+1}, ..., K_z$, and denote the resulting multicurve by $\boldsymbol{\zeta}''$. Notice that $\boldsymbol{\zeta}''$ belongs to $\mathcal{M}(n)$ and satisfies

$$\mathcal{S}_k(\boldsymbol{\zeta}'') \leq \mathcal{S}_k(\boldsymbol{\zeta}) - \mathcal{S}_k(\xi_v) - ... - \mathcal{S}_k(\xi_z) < \mathcal{S}_k(\boldsymbol{\zeta}) = \mathcal{S}_k(\boldsymbol{\gamma}),$$

which contradicts (2.6). ■

Lemma 2.20. *$\#B_{i,i} \leq 4(\ell_{\max}/\rho_{\text{inj}})^2 c + 16(\ell_{\max}/\rho_{\text{inj}})^3$, where c is the constant given by Lemma 2.19.*

Proof. By (2.11), (2.12), Lemma 2.18, and Lemma 2.19, we have

$$\begin{aligned}\#B_{i,i} &= \sum_{1\le h\le x\le a+b} \#\big(B_{i,i}\cap C_{h,x}\big)\\ &\le \sum_{1\le h<x\le a+b} \#C_{h,x} + \sum_{h=1}^{b} \#C_{a+h,a+h}\\ &\le (a+b)^2 c + (4\ell_{\max}/\rho_{\mathrm{inj}})^2 b\\ &\le 4(\ell_{\max}/\rho_{\mathrm{inj}})^2 c + 16(\ell_{\max}/\rho_{\mathrm{inj}})^3.\end{aligned}$$

■

Summing up, Lemmas 2.14, 2.16, 2.17, and 2.20 provide the following compactness result.

Proposition 2.21. *There exists $n_{\min}\ge n_{\mathrm{neg}}$ such that, for all integers $n\ge n_{\min}$, a minimizer of $\mathcal{S}_k$ over the space $\mathcal{M}(n)$ that does not have collapsed connected components belongs to $\mathcal{M}(n_{\min})$.* ■

Proof of Theorem 1.1. By Lemma 2.7, up to replacing M by a finite covering space and lifting the Lagrangian L to the tangent bundle of such covering space, for all integers $n\ge n_{\mathrm{neg}}$ the action functional $\mathcal{S}_k:\mathcal{M}(n)\to\mathbb{R}$ attains negative values. Let $\boldsymbol{\gamma}\in\mathcal{M}(n_{\min})$ be a multicurve without collapsed connected components such that

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \min_{\mathcal{M}(n_{\min})} \mathcal{S}_k < 0.$$

Proposition 2.21 implies that

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \min_{\mathcal{M}(n)} \mathcal{S}_k, \qquad \forall n\in\mathbb{N}. \tag{2.15}$$

Therefore, Lemmas 2.8 and 2.9 imply that the connected components of $\boldsymbol{\gamma}$ are embedded periodic orbits of the Lagrangian system of L with energy k. Since $\mathcal{S}_k(\boldsymbol{\gamma})<0$, at least one connected component of $\boldsymbol{\gamma}$, say γ_1, satisfies $\mathcal{S}_k(\gamma_1)<0$.

It only remains to show that the periodic orbit $\gamma_1:\mathbb{R}/\tau_1\mathbb{Z}\to M$ is a local minimizer of the free-period action functional $\mathcal{S}_k$ over the space of absolutely continuous periodic curves. We prove this by contradiction, assuming that there exists a sequence of absolutely continuous periodic curves $\gamma_{\alpha,1}:\mathbb{R}/\tau_{\alpha,1}\mathbb{Z}\to M$ such that

$$\begin{aligned}&\lim_{\alpha\to\infty}\tau_{\alpha,1}=\tau_1,\\ &\lim_{\alpha\to\infty}\max\big\{\mathrm{dist}(\gamma_{\alpha,1}(t/\tau_{\alpha,1}),\gamma_1(t/\tau_1))\;\big|\;t\in[0,1]\big\}=0,\\ &\mathcal{S}_k(\gamma_{\alpha,1})<\mathcal{S}_k(\gamma_1),\quad\forall\alpha\in\mathbb{N}.\end{aligned}$$

We fix an integer $h\in\mathbb{N}$ large enough so that, for all $\alpha\in\mathbb{N}$ sufficiently large and for all $t_0,t_1\in\mathbb{R}$ with $t_1-t_0<1/h$, we have $\mathrm{dist}(\gamma_{\alpha,1}(t_0),\gamma_{\alpha,1}(t_1))<\rho_{\mathrm{inj}}$. For all $j=0,...,h-1$, we set $t_{\alpha,j}:=\tau_{\alpha,1}j/h$, $q_{\alpha,j}:=\gamma_{\alpha,1}(t_{\alpha,j})$, $t_j:=\tau_1 j/h$, and $q_j:=\gamma_1(t_j)$. For each $j\in\mathbb{Z}/h\mathbb{Z}$, we remove from $\gamma_{\alpha,1}$ each portion $\gamma_{\alpha,1}|_{[t_{\alpha,j},t_{\alpha,j+1}]}$, and we glue in the unique free-time local minimizer with energy k joining $q_{\alpha,j}$ and $q_{\alpha,j+1}$. We still denote the resulting curve by $\gamma_{\alpha,1}:\mathbb{R}/\tau_{\alpha,1}\mathbb{Z}\to M$. Since we replaced portions of the original curve with unique free-time local minimizers with energy k, we still have $\mathcal{S}_k(\gamma_{\alpha,1})<\mathcal{S}_k(\gamma_1)$ for all $\alpha\in\mathbb{N}$. Moreover, since $\gamma_{\alpha,1}$ is now a piecewise solution of the Euler-Lagrange equation of L, and since its vertices

satisfy $q_{\alpha,j} \to q_j$ as $\alpha \to \infty$, for all $t \in [0,1]$ we have $\gamma_{\alpha,1}(\tau_{\alpha,1}t) \to \gamma_1(\tau_1 t)$ as $\alpha \to \infty$, and for all but finitely many $t \in [0,1]$ we have $\dot\gamma_{\alpha,1}(\tau_{\alpha,1}t) \to \dot\gamma_1(\tau_1 t)$ as $\alpha \to \infty$. Since γ_1 is a smoothly embedded curve, and since the property of being smoothly embedded is open in the C^1-topology, for all α large enough the curve $\gamma_{\alpha,1}$ is topologically embedded. Therefore, the multicurve $\boldsymbol{\gamma}_\alpha$ obtained by replacing γ_1 with $\gamma_{\alpha,1}$ in $\boldsymbol{\gamma}$, belongs to $\mathcal{M}(n)$ for some $n \in \mathbb{N}$, and satisfies

$$\mathcal{S}_k(\boldsymbol{\gamma}_\alpha) = \mathcal{S}_k(\boldsymbol{\gamma}) - \mathcal{S}_k(\gamma_1) + \mathcal{S}_k(\gamma_{\alpha,1}) < \mathcal{S}_k(\boldsymbol{\gamma}).$$

This contradicts (2.15). ∎

3. Free-period discretizations

3.1. **Unique action minimizers.** We work under the assumptions of Section 2.1: we consider a closed manifold M of arbitrary dimension, and a Tonelli Lagrangian $L : \mathrm{T}M \to \mathbb{R}$. We fix an energy value $k \in \mathbb{R}$. Since we are only interested in the Euler-Lagrange dynamics on $E^{-1}(k)$, up to modifying L far from $E^{-1}(-\infty, k]$ we can always assume that L is quadratic at infinity, that is, $L(q,v) = \frac{1}{2} g_q(v,v)$ for all $(q,v) \in \mathrm{T}M$ outside a compact set of $\mathrm{T}M$. Here, g is some fixed Riemannian metric on M. We will denote by $\mathrm{dist} : M \times M \to [0,\infty)$ the Riemannian distance associated to g.

We consider the free-period action functional $\mathcal{S}_k$ defined on the space of $W^{1,2}$ periodic curves on M of any period. Formally, such functional has the form

$$\mathcal{S}_k : W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty) \to \mathbb{R},$$
$$\mathcal{S}_k(\Gamma,\tau) = \tau \int_0^1 \Big[L(\Gamma(s), \Gamma'(s)/\tau) + k \Big] \,\mathrm{d}s = \int_0^\tau L(\gamma(t), \dot\gamma(t)) \,\mathrm{d}t + \tau k,$$

where we have identified (Γ,τ) with the τ-periodic curve $\gamma(t) = \Gamma(t/\tau)$. In this setting, $\mathcal{S}_k$ is $C^{1,1}$, but in general not C^2 unless the restriction of the Lagrangian L to any fiber of $\mathrm{T}M$ is a polinomial of degree two (see [AS09, Proposition 3.2]). The aim of this section is to overcome this lack of regularity by restricting $\mathcal{S}_k$ to a suitable finite dimensional submanifold, in the spirit of Morse's broken geodesics approximation of path spaces [Mil63, Section 16]. In the simpler setting of the fixed-period action functional, this type of analysis has been carried over by the second author in [Maz11, Section 3] and [Maz12, Chapter 4].

In Section 2.1, we investigated the properties of curves that are unique free-time local minimizers. In this section, we will need the following slightly different notion. An absolutely continuous curve $\gamma : [0,\tau] \to M$ is called a **unique action minimizer** when, for every other absolutely continuous curve $\zeta : [0,\tau] \to M$ with $\zeta(0) = \gamma(0)$ and $\zeta(\tau) = \gamma(\tau)$, we have

$$\int_0^\tau L(\gamma(t), \dot\gamma(t)) \,\mathrm{d}t < \int_0^\tau L(\zeta(t), \dot\zeta(t)) \,\mathrm{d}t.$$

It is well known that every unique action minimizer γ is a smooth solution of the Euler-Lagrange equation (1.1). A **Jacobi vector field** along such a curve γ is a solution $\theta : [0,\tau] \to \gamma^*(\mathrm{T}M)$ of the linearized Euler-Lagrange equation $\mathcal{J}_\gamma \theta = 0$, where $\mathcal{J}_\gamma$ is the linear second-order differential operator

$$\mathcal{J}_\gamma \theta := \tfrac{\mathrm{d}}{\mathrm{d}t} \Big[L_{qv}(\gamma,\dot\gamma)\,\theta + L_{vv}(\gamma,\dot\gamma)\,\dot\theta \Big] - L_{qq}(\gamma,\dot\gamma)\,\theta - L_{vq}(\gamma,\dot\gamma)\,\dot\theta.$$

A unique action minimizer $\gamma : [0,\tau] \to M$ is said to be **non-degenerate** when there is no Jacobi vector field $\theta : [0,\tau] \to \gamma^*(\mathrm{T}M)$ vanishing at both endpoints $\theta(0)$ and $\theta(1)$.

As before, we denote by $\phi_L^t : \mathrm{T}M \to \mathrm{T}M$ the Euler Lagrange flow of L, and by $\pi : \mathrm{T}M \to M$ the projection map of the tangent bundle. For $\rho > 0$, we define the open neighborhood of the diagonal in $M \times M$

$$\Delta_\rho := \big\{(q_0,q_1) \in M \times M \;\big|\; \mathrm{dist}(q_0,q_1) < \rho\big\}.$$

The following proposition follows from the proofs of [Maz12, Theorems 4.1.1 and 4.1.2].

Proposition 3.1. *There exist $\rho = \rho(L) > 0$, $\epsilon = \epsilon(L) > 0$, and smooth vector fields $\nu^\pm : \Delta_\rho \times (0,\epsilon) \to \mathrm{T}M$ such that, for all $(q_0,q_1,\tau) \in \Delta_\rho \times (0,\epsilon)$, we have*

$$\nu^-(q_0,q_1,\tau) \in \mathrm{T}_{q_0}M,$$
$$\nu^+(q_0,q_1,\tau) \in \mathrm{T}_{q_1}M,$$
$$\phi_L^\tau\big(q_0,\nu^-(q_0,q_1,\tau)\big) = \big(q_1,\nu^+(q_0,q_1,\tau)\big).$$

The curve $\gamma_{q_0,q_1,\tau} : [0,\tau] \to M$ given by

$$\gamma_{q_0,q_1,\tau}(t) := \pi \circ \phi_L^t\big(q_0,\nu^-(q_0,q_1,\tau)\big) = \pi \circ \phi_L^{t-\tau}\big(q_1,\nu^+(q_0,q_1,\tau)\big)$$

is a non-degenerate unique action minimizer. ∎

We need to consider the space of smooth paths defined on any compact interval of the form $[0,\tau]$. We can see this space as the product $C^\infty([0,1];M) \times (0,\infty)$ by identifying any pair $(\Gamma,\tau) \in C^\infty([0,1];M) \times (0,\infty)$ with the smooth curve $\gamma : [0,\tau] \to M$ given by $\gamma(t) = \Gamma(t/\tau)$. We define the smooth map

$$\iota : \Delta_\rho \times (0,\epsilon) \to C^\infty([0,1];M) \times (0,\epsilon),$$
$$\iota(q_0,q_1,\tau) := \gamma_{q_0,q_1,\tau} = (\Gamma_{q_0,q_1,\tau},\tau).$$

Proposition 3.1 implies that this map is an injective immersion. Indeed, for all $(v_0,v_1) \in \mathrm{T}_{q_0}M \times \mathrm{T}_{q_1}M$, consider

$$(\Theta_{v_0,v_1},0) := \mathrm{d}\iota(q_0,q_1,\tau)\big[(v_0,v_1,0)\big].$$

The curve $\theta_{v_0,v_1}(t) := \Theta_{v_0,v_1}(t/\tau)$ is a Jacobi vector field along $\gamma_{q_0,q_1,\tau}$ satisfying $\theta_{v_0,v_1}(0) = v_0$ and $\theta_{v_0,v_1}(\tau) = v_1$. Notice that such a Jacobi vector field is unique: if θ' were another Jacobi vector field along $\gamma_{q_0,q_1,\tau}$ with $\theta'(0) = v_0$ and $\theta'(\tau) = v_1$, the difference $\theta_{v_0,v_1} - \theta'$ would be a Jacobi vector field that vanishes at the endpoints, and therefore vanishes identically due to the non-degeneracy of $\gamma_{q_0,q_1,\tau}$.

Consider now the Jacobi vector field $\partial_\tau\gamma_{q_0,q_1,\tau}(t)$ along $\gamma_{q_0,q_1,\tau}$, which satisfies

$$\partial_\tau\gamma_{q_0,q_1,\tau}(0) = 0,$$
$$\partial_\tau\gamma_{q_0,q_1,\tau}(\tau) = -\dot\gamma_{q_0,q_1,\tau}(\tau) = -\nu^+(q_0,q_1,\tau).$$

Such vector field enters the expression of the derivative of the map ι with respect to τ. Indeed, consider

$$(\Psi_{q_0,q_1,\tau},1) := \mathrm{d}\iota(q_0,q_1,\tau)\big[(0,0,1)\big].$$

The vector field $\Psi_{q_0,q_1,\tau}$ along $\Gamma_{q_0,q_1,\tau}$ is given by

$$\Psi_{q_0,q_1,\tau}(t) = \partial_\tau\Gamma_{q_0,q_1,\tau}(t) = (\partial_\tau\gamma_{q_0,q_1,\tau})(\tau t) + \dot\gamma_{q_0,q_1,\tau}(\tau t)\,t.$$

Hence, the associated vector field $\psi_{q_0,q_1,\tau}$ along $\gamma_{q_0,q_1,\tau}$ is given by

$$\psi_{q_0,q_1,\tau}(t) = \Psi_{q_0,q_1,\tau}(t/\tau) = \partial_\tau \gamma_{q_0,q_1,\tau}(t) + \dot\gamma_{q_0,q_1,\tau}(t)\, t/\tau \tag{3.1}$$

and satisfies $\psi_{q_0,q_1,\tau}(0) = \psi_{q_0,q_1,\tau}(\tau) = 0$. Notice that $\psi_{q_0,q_1,\tau}$ is not a Jacobi vector field. Indeed, it satisfies

$$\begin{aligned} \mathcal{J}_{\gamma_{q_0,q_1,\tau}}(\psi_{q_0,q_1,\tau}) &= \mathcal{J}_{\gamma_{q_0,q_1,\tau}}(\dot\gamma_{q_0,q_1,\tau}\, t/\tau) \\ &= \big(\tfrac{\mathrm{d}}{\mathrm{d}t} E_v(\gamma_{q_0,q_1,\tau}, \dot\gamma_{q_0,q_1,\tau}) + E_q(\gamma_{q_0,q_1,\tau}, \dot\gamma_{q_0,q_1,\tau})\big)/\tau. \end{aligned} \tag{3.2}$$

3.2. **The discrete free-period action functional.** For $h \in \mathbb{N}$, which will be chosen large enough later, consider the open neighborhood of the h-fold diagonal

$$\Delta_{h,\rho} := \big\{ \boldsymbol{q} = (q_0, q_1, ..., q_{h-1}) \in M^{\times h} \;\big|\; \mathrm{dist}(q_i, q_{i+1}) < \rho,\ \forall i \in \mathbb{Z}/h\mathbb{Z} \big\}.$$

We equip $\Delta_{h,\rho} \times (0,\epsilon)$ with the Riemannian metric $\langle\!\langle \cdot,\cdot \rangle\!\rangle_\cdot$ given by

$$\langle\!\langle (\boldsymbol{v},\sigma), (\boldsymbol{w},\mu) \rangle\!\rangle_{(\boldsymbol{q},\tau)} = h\sigma\mu + \sum_{i\in\mathbb{Z}/h\mathbb{Z}} \langle v_i, w_i \rangle_{q_i}, \tag{3.3}$$

where $\langle\cdot,\cdot\rangle_\cdot$ denotes the Riemannian metric g on the manifold M. We define the injective smooth map

$$\begin{aligned} \iota_h : \Delta_{h,\rho} \times (0,\epsilon) &\to W^{1,2}(\mathbb{T}, M) \times (0,\infty), \\ \iota_h(\boldsymbol{q},\tau) &:= \gamma_{\boldsymbol{q},\tau} = (\Gamma_{\boldsymbol{q},\tau}, \tau h), \end{aligned} \tag{3.4}$$

where

$$\gamma_{\boldsymbol{q},\tau}(i\tau + t) := \gamma_{q_i,q_{i+1},\tau}(t), \qquad \forall i \in \mathbb{Z}/h\mathbb{Z},\ \forall t \in [0,\tau].$$

For later purposes, let us compute the derivative of ι_h. For all $(\boldsymbol{v},\sigma) \in \mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho} \times \mathbb{R}$, we have

$$\mathrm{d}\iota_h(\boldsymbol{q},\tau)\big[(\boldsymbol{v},\sigma)\big] = \big(\Theta_{\boldsymbol{v}} + \sigma\, \Psi_{\boldsymbol{q},\tau}, h\sigma\big), \tag{3.5}$$

where $\Theta_{\boldsymbol{v}}$ and $\Psi_{\boldsymbol{q},\tau}$ are vector fields along $\Gamma_{\boldsymbol{q},\tau}$ defined as follows: if $\theta_{\boldsymbol{v}}(t) = \Theta_{\boldsymbol{v}}(t/(\tau h))$ and $\psi_{\boldsymbol{q},\tau}(t) = \Psi_{\boldsymbol{q},\tau}(t/(\tau h))$, then

$$\begin{aligned} \theta_{\boldsymbol{v}}(i\tau + t) &= \theta_{v_i,v_{i+1}}(t), \\ \psi_{\boldsymbol{q},\tau}(i\tau + t) &= \psi_{q_i,q_{i+1},\tau}(t). \end{aligned}$$

Notice that $\theta_{\boldsymbol{v}}$ is the unique continuous vector field along $\gamma_{\boldsymbol{q},\tau}$ whose restriction to any interval of the form $[i\tau,(i+1)\tau]$ is the Jacobi vector field with boundary conditions $\theta_{\boldsymbol{v}}(i\tau) = v_i$ and $\theta_{\boldsymbol{v}}((i+1)\tau) = v_{i+1}$. This implies that ι_h is an immersion. We define the **discrete free-period action functional** as the composition

$$S_k := \mathcal{S}_k \circ \iota_h : \Delta_{h,\rho} \times (0,\epsilon) \to \mathbb{R}. \tag{3.6}$$

More explicitely, S_k is given by

$$\begin{aligned} S_k(\boldsymbol{q},\tau) &= \tau h k + \int_0^{\tau h} L(\gamma_{\boldsymbol{q},\tau}(t), \dot\gamma_{\boldsymbol{q},\tau}(t))\,\mathrm{d}t \\ &= \tau h k + \sum_{i\in\mathbb{Z}/h\mathbb{Z}} \int_0^{\tau} L(\gamma_{q_i,q_{i+1},\tau}(t), \dot\gamma_{q_i,q_{i+1},\tau}(t))\,\mathrm{d}t \\ &= \tau h k + \sum_{i\in\mathbb{Z}/h\mathbb{Z}} \int_0^{\tau} L \circ \phi_L^t(q_i, \nu^-(q_i,q_{i+1},\tau))\,\mathrm{d}t. \end{aligned}$$

The last expression readily implies that S_k is a C^∞ function.

3.3. **The discrete least action principle.** The differential of the free-period action is given by

$$\begin{aligned} \mathrm{d}\mathcal{S}_k(\Gamma,\tau)\big[(\Theta,\sigma)\big] = &\int_0^\tau \Big[L_q(\gamma(t),\dot\gamma(t))\,\theta(t) + L_v(\gamma(t),\dot\gamma(t))\,\dot\theta(t)\Big]\,\mathrm{d}t \\ &+ \frac{\sigma}{\tau}\int_0^\tau \Big[k - E(\gamma(t),\dot\gamma(t))\Big]\,\mathrm{d}t, \end{aligned} \tag{3.7}$$

where $\theta(t) := \Theta(t/\tau)$. This, together with an integration by parts and a bootstrapping argument, readily implies that the critical points of $\mathcal{S}_k$ are precisely the periodic orbits of the Lagrangian system of L with energy k, see, e.g., [Abb13, Section 3.2]. Any given critical point of $\mathcal{S}_k$ is contained in the image of the map ι_h for a large enough integer h. In particular, its preimage under ι_h is a critical point of S_k. Let us now verify that all critical points of S_k are mapped under ι_h to critical points of $\mathcal{S}_k$. By (3.5) and (3.7), we compute

$$\begin{aligned} \mathrm{d}S_k(\boldsymbol{q},\tau)\big[(\boldsymbol{v},\sigma)\big] &= \mathrm{d}\mathcal{S}_k(\Gamma_{\boldsymbol{q},\tau},\tau h)\big[(\Theta_{\boldsymbol{v}} + \sigma\,\Psi_{\boldsymbol{q},\tau}, h\sigma)\big] \\ &= \int_0^{\tau h}\Big[L_q(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\theta_{\boldsymbol{v}} + L_v(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\dot\theta_{\boldsymbol{v}}\Big]\,\mathrm{d}t \\ &\quad + \sigma\int_0^{\tau h}\Big[L_q(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\psi_{\boldsymbol{q},\tau} + L_v(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\dot\psi_{\boldsymbol{q},\tau}\Big]\,\mathrm{d}t \\ &\quad + \frac{\sigma}{\tau}\int_0^{\tau h}\Big[k - E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\Big]\,\mathrm{d}t. \end{aligned}$$

We break the integrals in the three summands as the sum of integrals over intervals of the form $[i\tau,(i+1)\tau]$. The first summand becomes

$$\begin{aligned} &\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\int_0^\tau \Big[L_q(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\theta_{v_i,v_{i+1}} + L_v(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\dot\theta_{v_i,v_{i+1}}\Big]\,\mathrm{d}t \\ &\quad= \sum_{i\in\mathbb{Z}/h\mathbb{Z}} L_v(\gamma_{q_i,q_{i+1},\tau}(t),\dot\gamma_{q_i,q_{i+1},\tau}(t))\theta_{v_i,v_{i+1}}(t)\Big|_{t=0}^{t=\tau} \\ &\quad= \sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[L_v(q_i,\nu^+(q_{i-1},q_i,\tau)) - L_v(q_i,\nu^-(q_i,q_{i+1},\tau))\Big]v_i. \end{aligned}$$

The second summand vanishes, since

$$\begin{aligned} &\int_0^\tau \Big[L_q(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\psi_{q_i,q_{i+1},\tau} + L_v(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\dot\psi_{q_i,q_{i+1},\tau}\Big]\,\mathrm{d}t \\ &\quad= \sum_{i\in\mathbb{Z}/h\mathbb{Z}} L_v(\gamma_{q_i,q_{i+1},\tau}(t),\dot\gamma_{q_i,q_{i+1},\tau}(t))\,\psi_{q_i,q_{i+1},\tau}(t)\Big|_{t=0}^{t=\tau} \\ &\quad= \sum_{i\in\mathbb{Z}/h\mathbb{Z}} L_v(\gamma_{q_i,q_{i+1},\tau}(t),\dot\gamma_{q_i,q_{i+1},\tau}(t))\,\partial_\tau\gamma_{q_i,q_{i+1},\tau}(t)\Big|_{t=0}^{t=\tau} \\ &\qquad+ \sum_{i\in\mathbb{Z}/h\mathbb{Z}} L_v(\gamma_{q_i,q_{i+1},\tau}(t),\dot\gamma_{q_i,q_{i+1},\tau}(t))\,\dot\gamma_{q_i,q_{i+1},\tau}(t)\,t/\tau\Big|_{t=0}^{t=\tau} \\ &\quad= \sum_{i\in\mathbb{Z}/h\mathbb{Z}} L_v(q_{i+1},\nu^+(q_i,q_{i+1},\tau))(-\nu^+(q_i,q_{i+1},\tau) + \nu^+(q_i,q_{i+1},\tau)) \\ &\quad= 0. \end{aligned}$$

Since $E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})$ is constant on each interval of the form $[i\tau,(i+1)\tau]$, the third summand becomes

$$\sigma\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[k-E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\Big]=\sigma\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[k-E(q_i,\nu^-(q_i,q_{i+1},\tau))\Big].$$

Summing up, we have obtained

$$\begin{aligned}\mathrm{d}S_k(\boldsymbol{q},\tau)\big[(\boldsymbol{v},\sigma)\big]=&\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[L_v(q_i,\nu^+(q_{i-1},q_i,\tau))-L_v(q_i,\nu^-(q_i,q_{i+1},\tau))\Big]v_i\\&+\sigma\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[k-E(q_i,\nu^-(q_i,q_{i+1},\tau))\Big].\end{aligned}\tag{3.8}$$

In particular, the critical points of the free-period action functional S_k are precisely those $(\boldsymbol{q},\tau)$'s such that $\nu^+(q_{i-1},q_i,\tau)=\nu^-(q_i,q_{i+1},\tau)$ and $E(q_{i+1},\nu^+(q_i,q_{i+1},\tau)=k$ for all $i\in\mathbb{Z}/h\mathbb{Z}$, that is, such that $\gamma_{\boldsymbol{q},\tau}$ is a τh-periodic orbit with energy k.

3.4. **The Hessian of the free-period action functional.** We denote by $H_{\boldsymbol{q},\tau}$ the Hessian of the free-period action functional S_k at a critical point $(\boldsymbol{q},\tau)$. By differentiating equation (3.8) we find that the expression of $H_{\boldsymbol{q},\tau}$ is

$$\begin{aligned}&H_{\boldsymbol{q},\tau}((\boldsymbol{v},\sigma),(\boldsymbol{w},\mu))\\&\quad=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,v_i,\mathrm{d}\nu^+(q_{i-1},q_i,\tau)(w_{i-1},w_i,0)-\mathrm{d}\nu^-(q_i,q_{i+1},\tau)(w_i,w_{i+1},0)\rangle\\&\qquad+\sigma\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,w_i,\partial_\tau\nu^+(q_{i-1},q_i,\tau)-\partial_\tau\nu^-(q_i,q_{i+1},\tau)\rangle\\&\qquad+\mu\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,v_i,\partial_\tau\nu^+(q_{i-1},q_i,\tau)-\partial_\tau\nu^-(q_i,q_{i+1},\tau)\rangle\\&\qquad-\sigma\mu\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\partial_\tau\big(E(q_i,\nu^-(q_i,q_{i+1},\tau))\big).\end{aligned}$$

We denote by $h_{\boldsymbol{q},\tau}$ the restriction of the Hessian $H_{\boldsymbol{q},\tau}$ to the codimension-one vector subspace $\mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho}\times\{0\}$, which reads

$$\begin{aligned}h_{\boldsymbol{q},\tau}(\boldsymbol{v},\boldsymbol{w}):=H_{\boldsymbol{q},\tau}\big((\boldsymbol{v},0),(\boldsymbol{w},0)\big)&\\=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,v_i,\mathrm{d}\nu^+(q_{i-1},q_i,\tau)(w_{i-1},w_i,0)&\\-\mathrm{d}\nu^-(q_i,q_{i+1},\tau)(w_i,w_{i+1},0)\rangle&\\=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,v_i,\dot\theta_{\boldsymbol{w}}(i\tau^-)-\dot\theta_{\boldsymbol{w}}(i\tau^+)\rangle.&\end{aligned}$$

We recall that the **nullity** $\mathrm{nul}(h)$ of a symmetric bilinear form h is the dimension of its kernel, whereas the **index** $\mathrm{ind}(h)$ is the maximal dimension of a vector subspace of its domain over which h is negative definite. In the usual terminology of Morse theory, $\mathrm{ind}(H_{\boldsymbol{q},\tau})$ and $\mathrm{nul}(H_{\boldsymbol{q},\tau})$ are the Morse index and the nullity of the functional S_k at the critical point $(\boldsymbol{q},\tau)$.

A vector $\boldsymbol{v}\in\mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho}$ belongs to $\ker(h_{\boldsymbol{q},\tau})$ if and only if $\dot\theta_{\boldsymbol{v}}(i\tau^-)=\dot\theta_{\boldsymbol{v}}(i\tau^+)$ for all $i\in\mathbb{Z}/h\mathbb{Z}$, that is, if and only if $\theta_{\boldsymbol{v}}$ is a smooth $h\tau$-periodic Jacobi vector field,

thus satisfying

$$\begin{aligned} \mathrm{d}\phi_L^t(\gamma_{\boldsymbol{q},\tau}(0),\dot\gamma_{\boldsymbol{q},\tau}(0))\big[(\theta_{\boldsymbol{v}}(0),\dot\theta_{\boldsymbol{v}}(0))\big] &= (\theta_{\boldsymbol{v}}(t),\dot\theta_{\boldsymbol{v}}(t)), \qquad \forall t\in\mathbb{R}, \\ (\theta_{\boldsymbol{v}}(0),\dot\theta_{\boldsymbol{v}}(0)) &= (\theta_{\boldsymbol{v}}(h\tau),\dot\theta_{\boldsymbol{v}}(h\tau)). \end{aligned}$$

This shows that the linear map $\boldsymbol{v}\mapsto(\theta_{\boldsymbol{v}}(0),\dot\theta_{\boldsymbol{v}}(0))$ gives an isomorphism between $\ker(h_{\boldsymbol{q},\tau})$ and the eigenspace of the linear map $P:=\mathrm{d}\phi_L^{h\tau}(\gamma_{\boldsymbol{q},\tau}(0),\dot\gamma_{\boldsymbol{q},\tau}(0))$ corresponding to the eigenvalue 1. In particular,

$$\mathrm{nul}(h_{\boldsymbol{q},\tau})=\dim\ker(P-I). \tag{3.9}$$

A vector $(\boldsymbol{v},\sigma)$ belongs to $\ker(H_{\boldsymbol{q},\tau})$ if and only if

$$\mathrm{d}\nu^+(q_{i-1},q_i,\tau)\big[(v_{i-1},v_i,\sigma)\big]=\mathrm{d}\nu^-(q_i,q_{i+1},\tau)\big[(v_i,v_{i+1},\sigma)\big] \tag{3.10}$$

and

$$\begin{aligned} \sum_{i\in\mathbb{Z}/h\mathbb{Z}} &\big\langle L_{vv}\,v_i,\partial_\tau\nu^+(q_{i-1},q_i,\tau)-\partial_\tau\nu^-(q_i,q_{i+1},\tau)\big\rangle \\ &= \sigma\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\partial_\tau\big(E(q_i,\nu^-(q_i,q_{i+1},\tau))\big). \end{aligned} \tag{3.11}$$

Conditions (3.10) and (3.11) can be conveniently rephrased by employing the vector fields $\theta_{\boldsymbol{v}}$ and $\psi_{\boldsymbol{q},\tau}$.

Lemma 3.2. *The kernel $\ker(H_{\boldsymbol{q},\tau})$ is isomorphic to the vector space of pairs (ξ,σ), where ξ is a smooth $h\tau$-periodic vector fields along $\gamma_{\boldsymbol{q},\tau}$ such that*

$$\mathcal{J}_{\gamma_{\boldsymbol{q},\tau}}\xi=\big(\tfrac{\mathrm{d}}{\mathrm{d}t}E_v(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})+E_q(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big)\sigma/\tau, \tag{3.12}$$

and $\sigma\in\mathbb{R}$ satisfies

$$\sigma\int_0^{h\tau}\langle L_{vv}(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\dot\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau}\rangle\,\mathrm{d}t=\tau\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(\xi,\dot\xi)\big]\,\mathrm{d}t. \tag{3.13}$$

The isomorphism is given by $(\boldsymbol{v},\sigma)\mapsto(\xi:=\theta_{\boldsymbol{v}}+\sigma\psi_{\boldsymbol{q},\tau},\sigma)$.

Remark 3.3. This lemma, together with the results in [AMP15, Appendix A], shows that the map $\mathrm{d}\iota_h(\boldsymbol{q},\tau)$ restricts to an isomorphism from $\ker(H_{\boldsymbol{q},\tau})$ to the kernel of the Hessian of the free-period action functional $\mathcal{S}_k$ at $(\gamma_{\boldsymbol{q},\tau},h\tau)$. Notice that there is a unique σ satisfying (3.13) unless $\gamma_{\boldsymbol{q},\tau}$ is a stationary curve. ■

Proof of Lemma 3.2. By the definition of the map ν^+, we have

$$\begin{aligned} \mathrm{d}\nu^+(q_{i-1},q_i,\tau)(v_{i-1},v_i,\sigma) &= \dot\theta_{\boldsymbol{v}}(i\tau^-)+\sigma\left(\partial_\tau\dot\gamma_{\boldsymbol{q},\tau}(i\tau^-)+\ddot\gamma_{\boldsymbol{q},\tau}(i\tau^-)\right) \\ &= \dot\theta_{\boldsymbol{v}}(i\tau^-)+\sigma\left(\dot\psi_{\boldsymbol{q},\tau}(i\tau^-)-\dot\gamma_{\boldsymbol{q},\tau}(i\tau)/\tau\right), \end{aligned}$$

and analogously

$$\begin{aligned} \mathrm{d}\nu^-(q_i,q_{i+1},\tau)(v_i,v_{i+1},\sigma) &= \dot\theta_{\boldsymbol{v}}(i\tau^+)+\sigma\partial_\tau\dot\gamma_{\boldsymbol{q},\tau}(i\tau^+) \\ &= \dot\theta_{\boldsymbol{v}}(i\tau^+)+\sigma\left(\dot\psi_{\boldsymbol{q},\tau}(i\tau^+)-\dot\gamma_{\boldsymbol{q},\tau}(i\tau)/\tau\right). \end{aligned}$$

We employ these two equations to rewrite (3.10) as

$$\dot\theta_{\boldsymbol{v}}(i\tau^+)+\sigma\dot\psi_{\boldsymbol{q},\tau}(i\tau^+)=\dot\theta_{\boldsymbol{v}}(i\tau^-)+\sigma\dot\psi_{\boldsymbol{q},\tau}(i\tau^-).$$

Namely, for each $(\boldsymbol{v},\sigma)\in\ker(H_{\boldsymbol{q},\tau})$, the $h\tau$-periodic vector field $\xi:=\theta_{\boldsymbol{v}}+\sigma\psi_{\boldsymbol{q},\tau}$ is C^1 and, by (3.2), satisfies

$$\mathcal{J}_{\gamma_{\boldsymbol{q},\tau}}(\xi)=\underbrace{\mathcal{J}_{\gamma_{\boldsymbol{q},\tau}}(\theta_{\boldsymbol{v}})}_{=0}+\sigma\mathcal{J}_{\gamma_{\boldsymbol{q},\tau}}(\psi_{\boldsymbol{q},\tau})=\big(\tfrac{\mathrm{d}}{\mathrm{d}t}E_v(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})+E_q(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big)\sigma/\tau.$$

In particular, ξ is C^∞.

Notice that, for all $\boldsymbol{v}\in\mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho}$, we have

$$\begin{aligned}\int_0^{h\tau}&\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(\theta_{\boldsymbol{v}},\dot\theta_{\boldsymbol{v}})\big]\,\mathrm{d}t\\&=\tau\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big[E_q(q_i,\nu^-(q_i,q_{i+1},\tau))v_i\\&\qquad+E_v(q_i,\nu^-(q_i,q_{i+1},\tau))\mathrm{d}\nu^-(q_i,q_{i+1},\tau)\big[(v_i,v_{i+1},0)\big]\Big]\\&=-\tau\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\langle L_{vv}\,v_i,\partial_\tau\nu^+(q_{i-1},q_i,\tau)-\partial_\tau\nu^-(q_i,q_{i+1},\tau)\rangle.\end{aligned}$$

Moreover

$$\begin{aligned}\int_0^{h\tau}&\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(\psi_{\boldsymbol{q},\tau},\dot\psi_{\boldsymbol{q},\tau})\big]\,\mathrm{d}t\\&=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\int_0^\tau\Big(\mathrm{d}E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\big[(\partial_\tau\gamma_{q_i,q_{i+1},\tau},\partial_\tau\dot\gamma_{q_i,q_{i+1},\tau})\big]\\&\qquad+\underbrace{\mathrm{d}E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\big[(\dot\gamma_{q_i,q_{i+1},\tau}t/\tau,\ddot\gamma_{q_i,q_{i+1},\tau}t/\tau)\big]}_{=0}\\&\qquad+\mathrm{d}E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\big[(0,\dot\gamma_{q_i,q_{i+1},\tau}/\tau)\big]\,\mathrm{d}t\Big)\\&=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\int_0^\tau\partial_\tau\big(E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\big)\,\mathrm{d}t\\&\quad+\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(0,\dot\gamma_{\boldsymbol{q},\tau})\big]\,\mathrm{d}t\\&=\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\Big(\partial_\tau\int_0^\tau E(\gamma_{q_i,q_{i+1},\tau},\dot\gamma_{q_i,q_{i+1},\tau})\,\mathrm{d}t-k\Big)\\&\quad+\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(0,\dot\gamma_{\boldsymbol{q},\tau})\big]\,\mathrm{d}t\\&=\tau\sum_{i\in\mathbb{Z}/h\mathbb{Z}}\partial_\tau\big(E(q_i,\nu^-(q_i,q_{i+1},\tau))\big)+\int_0^{h\tau}\langle L_{vv}\dot\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau}\rangle\,\mathrm{d}t.\end{aligned}$$

We plug these last two computations in equation (3.11), and infer that, for all $(\boldsymbol{v},\sigma)\in\ker(H_{\boldsymbol{q},\tau})$,

$$\begin{aligned}-\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(\theta_{\boldsymbol{v}},\dot\theta_{\boldsymbol{v}})\big]\,\mathrm{d}t=\sigma\Big(&\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau})\big[(\psi_{\boldsymbol{q},\tau},\dot\psi_{\boldsymbol{q},\tau})\big]\,\mathrm{d}t\\&-\int_0^{h\tau}\langle L_{vv}\dot\gamma_{\boldsymbol{q},\tau},\dot\gamma_{\boldsymbol{q},\tau}\rangle\,\mathrm{d}t\Big).\end{aligned}$$

Namely, σ satisfies (3.13) for $\xi := \theta_{\boldsymbol{v}} + \sigma\psi_{\boldsymbol{q},\tau}$.

On the other hand, if ξ is an $h\tau$-periodic vector field along $\gamma_{\boldsymbol{q},\tau}$ that satisfies equation (3.12) and $\sigma \in \mathbb{R}$ satisfies (3.13), the difference $\xi - \sigma\psi_{\boldsymbol{q},\tau}$ is an $h\tau$-periodic continuous and piecewise-smooth Jacobi vector field, smooth on each interval of the form $[i\tau,(i+1)\tau]$. Therefore, there exists $\boldsymbol{v} \in \mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho}$ such that $\theta_{\boldsymbol{v}} = \xi - \sigma\psi_{\boldsymbol{q},\tau}$. ■

Assume that the critical point $(\boldsymbol{q},\tau)$ of S_k is such that

$$\mathrm{dist}(\gamma_{\boldsymbol{q},\tau}(t_0),\gamma_{\boldsymbol{q},\tau}(t_1)) < \rho, \qquad \forall t_0,t_1 \in \mathbb{R} \text{ with } |t_1 - t_0| \le \tau.$$

In particular, $(\boldsymbol{q},\tau)$ belongs to a circle of critical points $(\boldsymbol{q}(s),\tau) \in \Delta_{h,\rho}$, for $s \in \mathbb{R}/h\tau\mathbb{Z}$, associated to the curves $\gamma_{\boldsymbol{q}(s),\tau} := \gamma_{\boldsymbol{q},\tau}(s+\cdot\,)$.

Lemma 3.4. *The indices* $\mathrm{ind}(h_{\boldsymbol{q}(s),\tau})$, $\mathrm{nul}(h_{\boldsymbol{q}(s),\tau})$, $\mathrm{ind}(H_{\boldsymbol{q}(s),\tau})$, *and* $\mathrm{nul}(H_{\boldsymbol{q}(s),\tau})$ *are independent of* $s \in \mathbb{R}/h\tau\mathbb{Z}$.

Proof. For each $s \in \mathbb{R}/h\tau\mathbb{Z}$ we set $P_s := \mathrm{d}\phi_L^{h\tau}(\gamma_{\boldsymbol{q},\tau}(s),\dot\gamma_{\boldsymbol{q},\tau}(s))$, so that

$$\mathrm{nul}(h_{\boldsymbol{q}(s),\tau}) = \dim\ker(P_s).$$

The linear map $(\theta_{\boldsymbol{v}}(0),\dot\theta_{\boldsymbol{v}}(0)) \mapsto (\theta_{\boldsymbol{v}}(s),\dot\theta_{\boldsymbol{v}}(s))$ is an isomorphism from $\ker(P_0)$ to $\ker(P_s)$, and in particular the function $s \mapsto \mathrm{nul}(h_{\boldsymbol{q}(s),\tau})$ is constant on $\mathbb{R}/h\tau\mathbb{Z}$.

The fact that the function $s \mapsto \mathrm{nul}(H_{\boldsymbol{q}(s),\tau})$ is constant on $\mathbb{R}/h\tau\mathbb{Z}$ is implied by Lemma 3.2. Indeed, assume that ξ and σ satisfy (3.12) and (3.13). For each $s \in \mathbb{R}/h\tau\mathbb{Z}$, the shifted vector field $\xi_s := \xi(s+\cdot\,)$ satisfies

$$\mathcal{J}_{\gamma_{\boldsymbol{q}(s),\tau}}\xi_s = \big(\tfrac{\mathrm{d}}{\mathrm{d}t}E_v(\gamma_{\boldsymbol{q}(s),\tau},\dot\gamma_{\boldsymbol{q}(s),\tau}) + E_q(\gamma_{\boldsymbol{q}(s),\tau},\dot\gamma_{\boldsymbol{q}(s),\tau})\big)\sigma/\tau,$$

$$\sigma\int_0^{h\tau}\langle L_{vv}(\gamma_{\boldsymbol{q}(s),\tau},\dot\gamma_{\boldsymbol{q}(s),\tau})\dot\gamma_{\boldsymbol{q}(s),\tau},\dot\gamma_{\boldsymbol{q}(s),\tau}\rangle\,\mathrm{d}t = \tau\int_0^{h\tau}\mathrm{d}E(\gamma_{\boldsymbol{q}(s),\tau},\dot\gamma_{\boldsymbol{q}(s),\tau})\big[(\xi_s,\dot\xi_s)\big]\,\mathrm{d}t.$$

Let A_s be the self-adjoint operator on $\mathrm{T}_{\boldsymbol{q}(s)}\Delta_{h,\rho}\times\mathbb{R}$ associated to $H_{\boldsymbol{q}(s),\tau}$, i.e.,

$$H_{\boldsymbol{q}(s),\tau}((\boldsymbol{v},\sigma),(\boldsymbol{w},\mu)) = \langle\!\langle A_s(\boldsymbol{v},\sigma),(\boldsymbol{w},\mu)\rangle\!\rangle_{(\boldsymbol{q}(s),\tau)}.$$

Since the eigenvalues depend continuously on the operator, there exist continuous functions $\lambda_i : \mathbb{R}/h\tau\mathbb{Z} \to \mathbb{R}$, for $i = 1,...,h\dim(M)$, such that the eigenvalues of each A_s are precisely the numbers $\lambda_i(s)$, repeated with their multiplicity. Since the nullity $\mathrm{nul}(H_{\boldsymbol{q}(s),\tau}) = \#\{i \mid \lambda_i(s) = 0\}$ is constant in s, we conclude that the index $\mathrm{ind}(H_{\boldsymbol{q}(s),\tau}) = \#\{i \mid \lambda_i(s) < 0\}$ is constant in s as well. An entirely analogous argument shows that $\mathrm{ind}(h_{\boldsymbol{q}(s),\tau})$ is constant is s as well. ■

3.5. **The iteration map.** The classical m-th **iteration map** on the free loop space sends any loop to its m-fold covering. In our finite dimensional setting, the analogous map $\psi^m : \Delta_{h,\rho}\times(0,\epsilon) \hookrightarrow \Delta_{mh,\rho}\times(0,\epsilon)$ is given by

$$\psi^m(\boldsymbol{q},\tau) = (\boldsymbol{q}^m,\tau),$$

where $\boldsymbol{q}^m = (\boldsymbol{q},\boldsymbol{q},...,\boldsymbol{q})$ denotes the m-fold diagonal vector. This is essentially a "linear" embedding (it would be a linear map in the formal sense if M were a Euclidean space instead of a closed manifold), and its differential is given by

$$\mathrm{d}\psi^m(\boldsymbol{q},\tau)\big[(\boldsymbol{v},\sigma)\big] = (\boldsymbol{v}^m,\sigma).$$

With a slight abuse of notation, we will use the symbol S_k to denote the free-period action functional on both spaces $\Delta_{h,\rho}\times(0,\epsilon)$ and $\Delta_{mh,\rho}\times(0,\epsilon)$. For all $(\boldsymbol{q},\tau) \in$

$\Delta_{h,\rho} \times (0,\epsilon)$ the gradient $(\boldsymbol{w},\mu) := \nabla S_k(\boldsymbol{q},\tau)$ with respect to the Riemannian metric (3.3) is given by

$$w_i = L_v(q_i,\nu^+(q_{i-1},q_i,\tau)) - L_v(q_i,\nu^-(q_i,q_{i+1},\tau)), \qquad \forall i \in \mathbb{Z}/h\mathbb{Z},$$
$$\mu = \frac{1}{h}\sum_{i\in\mathbb{Z}/h\mathbb{Z}} \Big[k - E(q_i,\nu^-(q_i,q_{i+1},\tau))\Big]$$

Analogously, the gradient $(\boldsymbol{w}',\mu') := \nabla S_k(\boldsymbol{q}^m,\tau)$ is given by

$$w'_i = w_{i \bmod h}, \qquad \forall i \in \mathbb{Z}/mh\mathbb{Z},$$
$$\mu' = \frac{1}{mh}\sum_{i\in\mathbb{Z}/mh\mathbb{Z}} \Big[k - E(q_i,\nu^-(q_i,q_{i+1},\tau))\Big] = \mu.$$

This proves the following.

Lemma 3.5. *For all $(\boldsymbol{q},\tau) \in \Delta_{h,\rho}\times(0,h\tau)$ we have*

$$\mathrm{d}\psi^m(\boldsymbol{q},\tau)\nabla S_k(\boldsymbol{q},\tau) = \nabla S_k(\psi^m(\boldsymbol{q},\tau)). \qquad \blacksquare$$

In order to simplify the notation, for all $m \in \mathbb{N}$ we abbreviate the Hessian bilinear forms as

$$h_m := h_{\boldsymbol{q}^m,\tau}, \qquad H_m := H_{\boldsymbol{q}^m,\tau}. \tag{3.14}$$

Lemma 3.6.

(i) *The function $m \mapsto \mathrm{nul}(H_m) - \mathrm{nul}(h_m)$ is constant on $\mathbb{N}$, and takes values in $\{-1,0,1\}$;*

(ii) *The function $m \mapsto \mathrm{ind}(H_m) - \mathrm{ind}(h_m)$ is constant on*

$$\mathbb{M} := \{m \in \mathbb{N} \mid \mathrm{nul}(h_1) = \mathrm{nul}(h_m)\}.$$

(iii) *The function $m \mapsto \mathrm{ind}(h_m)$ grows linearly, and is bounded if and only if it vanishes identically.*

Proof. We set

$$\mathbb{V}_m := \mathrm{T}_{\boldsymbol{q}^m}\Delta_{mh,\rho} \times \{0\},$$
$$\mathbb{K}_m := \ker(H_m) \cap \mathbb{V}_m = \ker(H_m) \cap \ker(h_m).$$

Notice that $\mathbb{K}_m$ is a vector subspace of $\ker(h_m)$ of codimension at most one, as well as a vector subspace of $\ker(H_m)$ of codimension at most one. Therefore

$$\mathrm{nul}(H_m) - \mathrm{nul}(h_m) \in \{-1,0,1\}.$$

We define the linear map

$$\omega^m : \mathrm{T}_{\boldsymbol{q}^m}(\Delta_{mh,\rho}) \times \mathbb{R} \to \mathrm{T}_{\boldsymbol{q}}(\Delta_{h,\rho}) \times \mathbb{R},$$
$$\omega^m(\boldsymbol{w},\mu) := (\boldsymbol{w}', m\mu),$$

where $w'_i = w_i + w_{i+h} + w_{i+2h} + \ldots + w_{i+(m-1)h}$ for all $i \in \mathbb{Z}/h\mathbb{Z}$. This map is the adjoint of $\psi^m_* := \mathrm{d}\psi^m(\boldsymbol{q},\tau)$ with respect to the Hessian of the free-period action functionals, in the sense that

$$H_m(\psi^m_*(\boldsymbol{v},\sigma),(\boldsymbol{w},\mu)) = H_1((\boldsymbol{v},\sigma),\omega^m(\boldsymbol{w},\mu)).$$

This equation readily implies that

$$\psi_*^m \ker(H_1) \subseteq \ker(H_m),$$
$$\psi_*^m \ker(h_1) \subseteq \ker(h_m),$$
$$\psi_*^m \mathbb{K}_1 \subseteq \mathbb{K}_m.$$

We recall that ψ_*^m is injective, since the iteration map ψ^m is an embedding. If $(\boldsymbol{v},0) \in \ker(h_1) \setminus \mathbb{K}_1$, then $H_1((\boldsymbol{v},0),(0,1)) \neq 0$ and

$$H_m(\psi_*^m(\boldsymbol{v},0),(0,1)) = H_1((\boldsymbol{v},0),\omega^m(0,1)) = H_1((\boldsymbol{v},0),(0,m)) \neq 0,$$

hence $\psi_*^m(\boldsymbol{v},0) \in \ker(h_m) \setminus \mathbb{K}_m$. Therefore, we have

$$\mathrm{nul}(h_m) - \dim(\mathbb{K}_m) = \mathrm{nul}(h_1) - \dim(\mathbb{K}_1), \qquad \forall m \in \mathbb{N}. \tag{3.15}$$

If $(\boldsymbol{v},\sigma) \in \ker(H_1) \setminus \mathbb{K}_1$, then $\sigma \neq 0$ and $\psi_*^m(\boldsymbol{v},\sigma) = (\boldsymbol{v}^m,\sigma) \in \ker(H_m) \setminus \mathbb{K}_m$. Therefore, we have

$$\mathrm{nul}(H_m) - \dim(\mathbb{K}_m) = \mathrm{nul}(H_1) - \dim(\mathbb{K}_1), \qquad \forall m \in \mathbb{N}. \tag{3.16}$$

Equations (3.15) and (3.16) imply point (i) of the lemma.

As for point (ii), consider again the vector space $\mathbb{V}_m$ defined above, and its H_m-orthogonal

$$\mathbb{V}_m^\perp := \Big\{ (\boldsymbol{v},\sigma) \in \mathrm{T}_{\boldsymbol{q}^m}\Delta_{mh,\rho} \times \mathbb{R} \;\Big|\; H_m((\boldsymbol{v},\sigma),\cdot)|_{\mathbb{V}_m} = 0 \Big\}.$$

An elementary formula from linear algebra (see, e.g., [Maz15, Prop. A.3]) allows to relate the index of any symmetric bilinear form to the index of its restriction to a vector subspace. For the Hessian H_m and the subspace $\mathbb{V}_m$, such formula reads

$$\mathrm{ind}(H_m) = \mathrm{ind}(h_m) + \mathrm{ind}(H_m|_{\mathbb{V}_m^\perp \times \mathbb{V}_m^\perp}) + \dim(\mathbb{V}_m \cap \mathbb{V}_m^\perp) - \dim(\mathbb{K}_m).$$

Notice that

$$\mathbb{V}_m \cap \mathbb{V}_m^\perp = \Big\{ (\boldsymbol{v},0) \in \mathbb{V}_m \;\Big|\; H_m((\boldsymbol{v},0),\cdot))|_{\mathbb{V}_m} = 0 \Big\} = \ker(h_m)$$

and therefore, by the very definition of the set $\mathbb{M}$, the intersection $\mathbb{V}_m \cap \mathbb{V}_m^\perp$ is independent of the integer $m \in \mathbb{M}$. Equation (3.15) further implies that $\dim(\mathbb{K}_m) = \dim(\mathbb{K}_1)$ for all $m \in \mathbb{M}$. In order to prove point (ii) of the lemma, we are only left to show that the index $\mathrm{ind}(H_m|_{\mathbb{V}_m^\perp \times \mathbb{V}_m^\perp})$ is independent of $m \in \mathbb{M}$. For all $(\boldsymbol{v},\sigma) \in \mathbb{V}_1^\perp$ and $(\boldsymbol{w},0) \in \mathbb{V}_m$, we have

$$H_m(\psi_*^m(\boldsymbol{v},\sigma),(\boldsymbol{w},0)) = H_1((\boldsymbol{v},\sigma),\omega^m(\boldsymbol{w},0)) = H_1((\boldsymbol{v},\sigma),(\boldsymbol{w}',0)) = 0.$$

On the other hand, for all $(\boldsymbol{v},\sigma) \in \mathbb{V}_m^\perp$ and $(\boldsymbol{w},0) \in \mathbb{V}_1$, we have

$$H_1(\omega^m(\boldsymbol{v},\sigma),(\boldsymbol{w},0)) = H_m((\boldsymbol{v},\sigma),\psi_*^m(\boldsymbol{w},0)) = H_m((\boldsymbol{v},\sigma),(\boldsymbol{w}^m,0)) = 0.$$

Namely,

$$\psi_*^m \mathbb{V}_1^\perp \subseteq \mathbb{V}_m^\perp,$$
$$\omega^m \mathbb{V}_m^\perp \subseteq \mathbb{V}_1^\perp.$$

The kernel $\ker(h_m)$ is contained in the vector space $\mathbb{V}_m^\perp$ with codimension at most one. For all $m \in \mathbb{M}$, we have $\psi_*^m \ker(h_1) = \ker(h_m)$. If there exists $(\boldsymbol{v},\sigma) \in \mathbb{V}_1^\perp \setminus \ker(h_1)$, then $\psi_*^m(\boldsymbol{v},\sigma) \notin \mathbb{V}_m$, and therefore $\psi_*^m(\boldsymbol{v},\sigma) \in \mathbb{V}_m^\perp \setminus \ker(h_m)$. Analogously,

if there exists $(\boldsymbol{v},\sigma)\in \mathbb{V}_m^\perp\setminus\ker(h_m)$, then $\omega^m(\boldsymbol{v},\sigma)\notin\mathbb{V}_1$, and therefore $\omega^m(\boldsymbol{v},\sigma)\in \mathbb{V}_1^\perp\setminus\ker(h_1)$. This shows that $\psi_*^m\mathbb{V}_1^\perp=\mathbb{V}_m^\perp$. Since, for all $(\boldsymbol{v},\sigma)\in\mathbb{V}_1^\perp$, we have

$$H_m(\psi_*^m(\boldsymbol{v},\sigma),\psi_*^m(\boldsymbol{v},\sigma))=mH_1((\boldsymbol{v},\sigma),(\boldsymbol{v},\sigma)),$$

we conclude that $\operatorname{ind}(H_m|_{\mathbb{V}_m^\perp\times\mathbb{V}_m^\perp})=\operatorname{ind}(H_1|_{\mathbb{V}_1^\perp\times\mathbb{V}_1^\perp})$ for all $m\in\mathbb{M}$.

Now, assume that the function $m\mapsto\operatorname{ind}(h_m)$ does not vanish identically, and consider $m_0\in\mathbb{N}$ such that $\operatorname{ind}(h_{m_0})>0$. In particular, there exists a non-zero vector $\boldsymbol{v}\in\mathrm{T}_{\boldsymbol{q}^{m_0}}\Delta_{m_0h,\rho}$ such that $\delta_1:=h_{m_0}(\boldsymbol{v},\boldsymbol{v})<0$. We set

$$\begin{aligned}\delta_2:=&\langle L_{vv}\,v_0,\mathrm{d}\nu^+(q_{h-1},q_0,\tau)\big[(0,v_0,0)\big]\rangle\\&-\langle L_{vv}\,v_{m_0h-1},\mathrm{d}\nu^-(q_{h-1},q_0,\tau)\big[(v_{m_0h-1},0,0)\big]\rangle,\\ m_1:=&\big\lceil|\delta_2/\delta_1|\big\rceil.\end{aligned}$$

We consider an integer $m\geq m_0m_1+1$, and define the vectors

$$\begin{aligned}\boldsymbol{w}_i:=(\mathbf{0}^{i(m_0m_1+1)},\boldsymbol{v}^{m_1},\mathbf{0}^{m-i(m_0m_1+1)-m_0m_1})\in\mathrm{T}_{\boldsymbol{q}^m}\Delta_{mh,\rho},\\ \forall i=0,...,\big\lfloor\tfrac{m}{m_0m_1+1}\big\rfloor-1,\end{aligned}$$

where $\mathbf{0}=(0,...,0)$ is the origin in $\mathrm{T}_{\boldsymbol{q}}\Delta_{h,\rho}$. Notice that, for all $i\neq j$, we have

$$\begin{aligned}h_m(\boldsymbol{w}_i,\boldsymbol{w}_i)&=m_1\delta_1+\delta_2<0,\\ h_m(\boldsymbol{w}_i,\boldsymbol{w}_j)&=0.\end{aligned}$$

Therefore, the $\boldsymbol{w}_i$'s span an $\big\lfloor\frac{m}{m_0m_1+1}\big\rfloor$-dimensional vector subspace over which the Hessian h_m is negative definite. In particular

$$\operatorname{ind}(h_m)\geq\big\lfloor\tfrac{m}{m_0m_1+1}\big\rfloor,\qquad\forall m\in\mathbb{N}.$$ ■

The following lemma is the analog of a closed geodesics result due to Gromoll and Meyer [GM69, Lemma 1.2] in our finite dimensional setting for the free-period action functional. In the infinite dimensional setting, the result was established in [AMMP14, Lemma 1.2].

Lemma 3.7. *Let $(\boldsymbol{q},\tau)$ be a critical point of the discrete free-period action functional S_k. The set of positive integers admits a partition $\mathbb{N}_1\cup...\cup\mathbb{N}_r$, integers $m_1\in\mathbb{N}_1$, ..., $m_r\in\mathbb{N}_r$, and $\nu_1,...,\nu_r\in\{0,1,...,2\dim(M)+1\}$ such that m_j divides all the integers in $\mathbb{N}_j$, and $\operatorname{nul}(H_m)=\nu_j$ for all $m\in\mathbb{N}_j$.*

Proof. Thanks to Lemma 3.6(i), it is enough to provide a proof for the nullity of h_m instead of the nullity of H_m. The argument leading to (3.9) shows that

$$\operatorname{nul}(h_m)=\dim\ker(P^m-I),\qquad\forall m\in\mathbb{N}.\tag{3.17}$$

The geometric multiplicity of the eigenvalue 1 of a power matrix varies as

$$\dim\ker(P^m-I)=\sum_{\lambda\in\sqrt[m]{1}}\dim_{\mathbb{C}}\ker_{\mathbb{C}}(P-\lambda I),\tag{3.18}$$

see, e.g., [Maz15, Prop. A.1]. The remaining of the proof is a standard arithmetic argument. We denote by $\sigma_m(P)$ the set of those eigenvalues of P on the complex unit circle that are m-th roots of unity, and we introduce the following equivalence relation on the natural numbers: $m\sim m'$ if and only if $\sigma_m(P)=\sigma_{m'}(P)$. The subsets $\mathbb{N}_1,...,\mathbb{N}_r$ of the lemma will be the equivalence classes of this relation. If m_j is the minimum of $\mathbb{N}_j$, and $\sigma_{m_j}(P)=\{\exp(i2\pi p_1/q_1),...,\exp(i2\pi p_s/q_s)\}$, the

set $\mathbb{N}_j$ consists precisely of the common multiples of $q_1, ..., q_s$. In particular, every $m \in \mathbb{N}_j$ is divisible by m_j and, by (3.17) and (3.18), we conclude

$$\mathrm{nul}(h_m) = \sum_{\lambda \in \sigma_m(P)} \dim_{\mathbb{C}} \ker_{\mathbb{C}}(P - \lambda I) = \sum_{\lambda \in \sigma_{m_j}(P)} \dim_{\mathbb{C}} \ker_{\mathbb{C}}(P - \lambda I) = \mathrm{nul}(h_{m_j}).$$

■

4. Multiplicity of low energy periodic orbits

This section is devoted to the proof of Theorem 1.2, which will follow closely the one for the electromagnetic case in [AMMP14]. The essential difference in our general Tonelli case is that we need to employ the finite dimensional functional setting of Section 3 in all the arguments involving the Hessian of the free-period action functional.

4.1. **A property of high iterates of periodic orbits.** Let M be a closed manifold of arbitrary positive dimension, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian. We fix an energy value $k \in \mathbb{R}$. Up to modifying L outside an open neighborhood of $E^{-1}(-\infty, k]$, we can assume that $L(q, v) = \frac{1}{2} g_q(v, v)$ outside a compact set of $\mathrm{T}M$, where g is some Riemannian metric on M. We will adopt the notation of Section 3, and in particular we consider the positive constants $\rho = \rho(L)$ and $\epsilon = \epsilon(L)$ given by Proposition 3.1. Let $\gamma : \mathbb{R}/\sigma\mathbb{Z} \to M$ be a periodic orbit of the Lagrangian system of L with energy k. We choose an integer $h \geq 1/\epsilon$ large enough so that

$$\mathrm{dist}(\gamma(t_0), \gamma(t_1)) < \rho, \qquad \forall t_0, t_1 \in \mathbb{R} \text{ with } |t_0 - t_1| \leq \sigma/h. \tag{4.1}$$

Therefore, there exists a critical point $(\boldsymbol{q}, \tau)$ of the discrete free-period action functional $S_k : \Delta_{h,\rho} \times (0, \epsilon) \to \mathbb{R}$ such that $\sigma = h\tau$ and $\gamma = \gamma_{\boldsymbol{q},\tau}$. We denote by

$$K = \bigcup_{s \in \mathbb{R}/h\tau\mathbb{Z}} \{(\boldsymbol{q}(s), \tau)\},$$

the critical circle containing $(\boldsymbol{q}, \tau)$, where $(\boldsymbol{q}(s), \tau)$ is the critical point associated to the periodic orbit

$$\gamma_{\boldsymbol{q}(s),\tau} := \gamma_{\boldsymbol{q},\tau}(s + \cdot).$$

Let $c := S_k(\boldsymbol{q}, \tau)$. We assume that K is isolated in $\mathrm{crit}(S_k)$, and therefore admits an arbitrarily small connected open neighborhood $U \subset \Delta_{h,\rho} \times (0, \epsilon)$ such that the intersection $\{S_k < c\} \cap U$ has only finitely many connected components $U_1^-, ..., U_l^-$ (this follows from the more general statement that the local homology of the isolated critical circle K has finite rank). We can assume that

$$\partial U_i^- \cap K \neq \varnothing, \qquad \forall i = 1, ..., l. \tag{4.2}$$

Indeed, if this is not verified, we remove from U all the U_i^-'s whose boundary does not intersect K, which leaves us with a smaller connected open neighborhood of K that satisfies (4.2).

Lemma 4.1. $K \subset \partial U_i^- \cap ... \cap \partial U_l^-$.

Proof. For each $(\boldsymbol{q}', \tau') \in U_i^-$ and $s \in \mathbb{R}$, we set

$$\boldsymbol{q}'(s) := \big(\gamma_{\boldsymbol{q}',\tau'}(s), \gamma_{\boldsymbol{q}',\tau'}(s + \tau'), ..., \gamma_{\boldsymbol{q}',\tau'}(s + (h-1)\tau')\big).$$

By (4.1), all points $\boldsymbol{q}'(s)$ belong to $\Delta_{h,\rho}$ provided $(\boldsymbol{q}', \tau')$ is sufficiently close to K. Moreover, $S_k(\boldsymbol{q}'(s), \tau') \leq S_k(\boldsymbol{q}', \tau') < c$, that is,

$$(\boldsymbol{q}'(s), \tau') \in \{S_k < c\}. \tag{4.3}$$

By (4.2), there exists $(\boldsymbol{q}(s_0), \tau) \in \partial U_i^- \cap K$. If we take a sequence of points $(\boldsymbol{q}', \tau') \in U_i^-$ such that $(\boldsymbol{q}', \tau') \to (\boldsymbol{q}(s_0), \tau)$, we have $(\boldsymbol{q}'(s), \tau') \to (\boldsymbol{q}(s_0 + s), \tau)$ for all $s \in (0, h\tau)$. This, together with (4.3), implies that $K \subset \partial U_i^-$. ∎

Lemma 4.2. *For all $m \in \mathbb{N}$ large enough, the image under the iteration map $\psi^m(\{S_k < c\} \cap U)$ is contained in one connected component of the sublevel set $\{S_k < mc\}$.*

Proof. Consider two distinct connected components U_1 and U_2 of $\{S_k < c\} \cap U$ (if this latter intersection is connected we are already done). We denote by $B_r \subset \Delta_{h,\rho} \times (0, \epsilon)$ an open ball of radius $r > 0$ centered at $(\boldsymbol{q}, \tau) \in K$. By Lemma 4.1, the critical point $(\boldsymbol{q}, \tau)$ belongs to the intersection $\partial U_1 \cap \partial U_2$. Therefore, there exists a continuous path $(\boldsymbol{q}', \tau') : [0, 1] \to B_r$ such that $(\boldsymbol{q}'(0), \tau'(0)) \in U_1$ and $(\boldsymbol{q}'(1), \tau'(1)) \in U_2$. Consider the embedding ι_h defined in (3.4), and the continuous path

$$\iota_h \circ (\boldsymbol{q}', \tau') : [0, 1] \to W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty),$$
$$\iota_h \circ (\boldsymbol{q}'(u), \tau'(u)) = (\Gamma_{\boldsymbol{q}'(u), \tau'(u)}, h\tau'(u)).$$

By Bangert's trick of "pulling one loop at the time" (see [Ban80, pages 86-87], [Abb13, page 421], or [Maz14, Section 3.2]), for all integers m large enough there exists a continuous homotopy

$$F_r : [0, 1] \to W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, mh\epsilon), \qquad r \in [0, 1],$$

such that $F_0 = \iota_{mh} \circ \psi^m \circ (\boldsymbol{q}', \tau')$, $F_r(0) = F_0(0)$ and $F_r(1) = F_0(1)$ for all $r \in [0, 1]$, and $F_1([0, 1]) \subset \{\mathcal{S}_k < mc\}$. Moreover, up to reducing the radius r of the original ball (and consequently increasing the integer m necessary for Bangert's trick), we can make F_r take values in an arbitrarily small C^0-neighborhood of the m-th iterate of $\gamma_{\boldsymbol{q},\tau} = (\Gamma_{\boldsymbol{q},\tau}, h\tau)$. We write the homotopy as $F_r(u) = (\Gamma_r''(u), mh\tau_r''(u))$. We define another continuous homotopy

$$f_r : [-1, 1] \to \Delta_{mh,\rho} \times (0, \epsilon),$$
$$f_r(u) := (\boldsymbol{q}''(r, u), \tau_r''(u)),$$

where

$$\boldsymbol{q}''(r, u) := \big(\Gamma_r''(u)(0), \Gamma_r''(u)(\tfrac{1}{mh}), ..., \Gamma_r''(u)(\tfrac{mh-1}{mh})\big).$$

Notice that $f_0(u) = (\boldsymbol{q}'(u), \tau'(u))$, $f_r(-1) = f_0(-1)$ and $f_r(1) = f_0(1)$ for all $r \in [0, 1]$, and $S_k(f_1(u)) \leq \mathcal{S}_k(F_1(u)) < mc$. The existence of the continuous path f_1 implies the lemma. ∎

One of the main ingredients for the proof of Theorem 1.2 is the following statement about high iterates of periodic orbits. In the special case of Tonelli Lagrangians that restrict to polynomials of degree 2 in any fiber of the tangent bundle, an infinite dimensional version of this theorem was established in [AMMP14, Theorem 2.6]. A similar statement for closed Riemannian geodesics on surfaces was originally established by Bangert in [Ban80, Theorem 2].

Lemma 4.3. *Assume that, for all $m \in \mathbb{N}$, the point $(\boldsymbol{q}^m, \tau)$ belongs to an isolated critical circle of $S_k : \Delta_{mh,\rho} \times (0,\epsilon) \to \mathbb{R}$. Then, if m is large enough, there exists a connected open neighborhood $W \subset \Delta_{mh,\rho} \times (0,\epsilon)$ of the critical circle of $(\boldsymbol{q}^m, \tau)$ such that the inclusion map induces the injective map among connected components*

$$\pi_0(\{S_k < mc\}) \hookrightarrow \pi_0(\{S_k < mc\} \cup W). \tag{4.4}$$

Remark 4.4. This lemma can be phrased by saying that sufficiently high iterates of periodic orbits are not 1-dimensional mountain passes for the discrete free-period action functional. By applying the same techniques as in [MM15, Section 4], one can prove that, for any given periodic orbit and degree $d \geq 1$, sufficiently high iterates of the periodic orbit are not d-dimensional mountain passes for the discrete free-period action functional. Since the case $d = 1$ suffices for the proof of Theorem 1.2, we will only present this case. ■

Proof of Lemma 4.3. As in (3.14), we denote by H_m and h_m the Hessians of the discrete free-period and fixed-period action functionals at $(\boldsymbol{q}^m, \tau)$. By Lemma 3.6(iii), either $\mathrm{ind}(h_m) \to \infty$ as $m \to \infty$ or $\mathrm{ind}(h_m) = 0$ for all $m \in \mathbb{N}$.

In the former case, by Lemma 3.6(i), for all m large enough we have $\mathrm{ind}(H_m) \geq 2$. By Lemma 3.4, the discrete free-period action functional S_k have the same Morse index (larger than or equal to 2) and nullity at all the critical points belonging to the critical circle of $(\boldsymbol{q}^m, \tau)$. In particular, there exists arbitrarily small connected open neighborhoods $W \subset \Delta_{mh,\rho} \times (0,\epsilon)$ of the critical circle of $(\boldsymbol{q}^m, \tau)$ such that the intersection $\{S_k < mc\} \cap W$ is connected, and (4.4) follows.

Let us now deal with the second case in which $\mathrm{ind}(h_m) = 0$ for all $m \in \mathbb{N}$. We denote by K_m the critical circle of S_k containing $(\boldsymbol{q}^m, \tau)$. We consider the partition $\mathbb{N} = \mathbb{N}_1 \cup ... \cup \mathbb{N}_r$ given by Lemma 3.7, and the integers $m_i := \min \mathbb{N}_i$. By Lemmas 3.6(ii) and 3.7, we have $\mathrm{ind}(H_m) = \mathrm{ind}(H_{m_i})$ and $\mathrm{nul}(H_m) = \mathrm{nul}(H_{m_i})$ for all $m \in \mathbb{N}_i$. By Lemma 4.2 and the prior discussion, for each $i \in \{1, ..., r\}$ there exists a connected open neighborhood U of K_{m_i} such that, for all $m \in \mathbb{N}_i$ large enough, $\psi^{m/m_i}(\{S_k < m_i c\} \cap U)$ is contained in one connected component of the sublevel set $\{S_k < mc\}$.

Let $U' \subset U$ be a sufficiently small connected neighborhood of K_{m_i}. Since the iteration map $\psi^{m/m_i} : \Delta_{m_i h,\rho} \times (0,\epsilon) \hookrightarrow \Delta_{mh,\rho} \times (0,\epsilon)$ is an embedding, $\psi^{m/m_i}(U')$ admits a tubular neighborhood $W \subset \Delta_{mh,\rho} \times (0,\epsilon)$, which we identify with an open neighborhood of the zero-section of the normal bundle of $\psi^{m/m_i}(U')$ with projection $\pi : W \to \psi^{m/m_i}(U')$. We denote the points of W by (x, v), where $x \in \psi^{m/m_i}(U')$ is a point in the base and $v \in \pi^{-1}(x)$ is a point in the corresponding fiber. By Lemma 3.5, the gradient of S_k is tangent to the zero-section of W at all points $(x, 0) \in W$. Therefore, the restriction of S_k to any fiber $\pi^{-1}(x)$ has a critical point at the origin. We introduce the radial deformation $r_t : W \to W$, for $t \in [0, 1]$, given by $r_t(x, v) = (x, (1 - t)v)$. This is a deformation retraction of the neighborhood W onto the base $\psi^{m/m_i}(U')$. By Lemma 3.4, every critical point of S_k on the critical circle $\psi^{m/m_i}(K)$ has Morse index $\mathrm{ind}(H_m)$ and nullity $\mathrm{nul}(H_m)$. Since $\mathrm{ind}(H_m) = \mathrm{ind}(H_{m_i})$ and $\mathrm{nul}(H_m) = \mathrm{nul}(H_{m_i})$, up to shrinking U' we have that the restriction of the discrete free-period action functional $S_k : W \to \mathbb{R}$ to any fiber $\pi^{-1}(x)$ has a non-degenerate local minimum at the origin; up to shrinking W, such local minimum is a global minimum and we have $\frac{\mathrm{d}}{\mathrm{d}t} S_k \circ r_t \leq 0$. In particular, the deformation r_t preserves the intersection $\{S_k < mc\} \cap W$. Since

$r_1(\{S_k < mc\} \cap W) = \psi^{m/m_i}(U')$, we conclude that $\{S_k < mc\} \cap W$ is contained in one connected component of $\{S_k < mc\}$, which implies (4.4). ■

4.2. **The minimax scheme.** From now on, we will further assume that M is a closed surface, i.e.,

$$\dim(M) = 2.$$

Up to lifting L to the tangent bundle of the orientation double covering of M, we can assume that M is orientable. We recall that the Mañé critical value $c_{\mathrm{u}}(L)$ is defined as the minimum k such that the free-period action functional $\mathcal{S}_k$ is non-negative on the connected component of contractible periodic curves, or equivalently as the usual Mañé critical value of the lift of L to the tangent bundle of the universal covering of M. It is easy to see that $e_0(L) \leq c_{\mathrm{u}}(L) \leq c_0(L)$.

In order to prove Theorem 1.2, we employ the periodic orbit provided by Theorem 1.1: since such periodic orbit is a local minimum of the free-period action functional $\mathcal{S}_k : W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty) \to \mathbb{R}$, and since this functional is unbounded from below on every connected component when $k < c_{\mathrm{u}}(L)$, we can find a second periodic orbit by performing a suitable 1-dimensional minimax. The framework of such minimax is not entirely standard, since $\mathcal{S}_k$ might not satisfy the Palais-Smale condition. The argument for the case of fiberwise quadratic Lagrangians provided in [AMMP14, Sections 3.1-3.2] actually works for general Tonelli Lagrangians; indeed, such argument only requires that $\mathcal{S}_k$ is C^1. We will thus present the construction and the properties of the minimax scheme (Lemma 4.5 below) without proofs, which the reader can find in [AMMP14, Sections 3.1-3.2].

We fix, once for all, an energy value $k_0 \in (e_0(L), c_{\mathrm{u}}(L))$. Let $\gamma_{k_0} = (\Gamma_{k_0}, \tau_{k_0}) \in W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty)$ be a local minimizer of $\mathcal{S}_{k_0}$ with action $\mathcal{S}_{k_0}(\gamma_{k_0}) < 0$, whose existence is guaranteed by Theorem 1.1. Since the configuration space M is assumed to be an orientable surface, every iterate of γ_{k_0} is a local minimizer of $\mathcal{S}_{k_0}$ as well, see [AMP15, Lemma 4.1]. The free-period action functional satisfies the Palais-Smale condition on subsets of the form $W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times [\epsilon_0, \epsilon_1]$, where $0 < \epsilon_0 \leq \epsilon_1 < \infty$. Therefore, we can find ϵ_0, ϵ_1 and a sufficiently small open neighborhood $\mathcal{U} \subset W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times [\epsilon_0, \epsilon_1]$ of the critical circle C of γ_{k_0} such that

$$\begin{aligned} &\mathcal{U} \cap \mathrm{crit}(\mathcal{S}_{k_0}) = C, \\ &\inf_{\partial\mathcal{U}} \mathcal{S}_{k_0} > \mathcal{S}_{k_0}(\gamma_{k_0}). \end{aligned}$$

The free-period action functional $\mathcal{S}_{k_0}$ is unbounded from below in every connected component, for $k_0 < c_{\mathrm{u}}(L)$. In particular, there exists $\zeta \in W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty)$ in the same connected component as γ_{k_0} such that $\mathcal{S}_{k_0}(\zeta) < \mathcal{S}_{k_0}(\gamma_{k_0})$. For each $k \in \mathbb{R}$, we denote by $\mathcal{M}_k \subset W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times [\epsilon_0, \epsilon_1]$ the (possibly empty) closure of the set of local minimizers of $\mathcal{S}_k|_{\mathcal{U}}$. Notice that $\mathcal{M}_{k_0} = C$. As customary, we denote by $\gamma^n : \mathbb{R}/n\tau\mathbb{Z} \to M$ the n-th iterate of a closed curve $\gamma : \mathbb{R}/\tau\mathbb{Z} \to M$. For all $n \in \mathbb{N}$ and $k \in \mathbb{R}$, we denote by $\mathcal{P}(n,k)$ the set of continuous paths

$$p : [0,1] \to W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty)$$

such that $p(0) = \zeta^n$ and $p(1) = \mu^n$ for some $\mu \in \mathcal{M}_k$. The following statement is quoted from [AMMP14, Lemma 3.5].

Lemma 4.5. *There exists a full-measure subset J of a closed neighborhood of k_0 such that, for all $n \in \mathbb{N}$ and $k \in J$, the real number*

$$c(n,k) := \inf_{p \in \mathcal{P}(n,k)} \max_{s \in [0,1]} \mathcal{S}_k(p(s))$$

is a critical value of the free-period action functional $\mathcal{S}_k$, and

$$\lim_{n \to \infty} c(n,k) = -\infty.$$

For every neighborhood $\mathcal{V}$ of $\mathrm{crit}(\mathcal{S}_k) \cap \mathcal{S}_k^{-1}(c(n,k))$ there exists $p \in \mathcal{P}(n,k)$ such that $p([0,1]) \subset \{\mathcal{S}_k < c(n,k)\} \cup \mathcal{V}$. ∎

Proof of Theorem 1.2. We prove the theorem by contradiction. We fix, once for all, an energy value $k_0 \in (e_0(L), c_{\mathrm{u}}(L))$ and the corresponding set $J \subset (e_0(L), c_{\mathrm{u}}(L))$ given by Lemma 4.5. It is enough to prove that, for each $k \in J$, the family of critical points of $\mathcal{S}_k$ with critical values in $\{c(n,k) \,|\, n \in \mathbb{N}\}$ contains infinitely many (geometrically distinct) periodic orbits. Assume by contradiction that this does not hold. Therefore, there exist finitely many critical points $\zeta_1, ..., \zeta_r$ of $\mathcal{S}_k$ such that, for all $n \in \mathbb{N}$, all the critical points of $\mathcal{S}_k$ with critical values $c(n,k)$ are iterates of some periodic orbits among $\zeta_1, ..., \zeta_r$.

We consider the finite dimensional setting of Section 3. For an integer $h \in \mathbb{N}$, we introduce the open subset $\mathcal{W} \subset W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$ given by those periodic curves $\gamma = (\Gamma, \tau) \in W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$ such that

$$\mathrm{dist}(\gamma(t_0), \gamma(t_1)) < \rho, \qquad \forall t_0, t_1 \in \mathbb{R} \text{ with } |t_0 - t_1| \leq \tau/h.$$

We define a homotopy $h_s : \mathcal{W} \to W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$, for $s \in [0,1]$, as follows. For each $s \in [0,1]$ and $\gamma = (\Gamma, \tau) \in \mathcal{W}$, we set

$$q_i := \gamma\big(\tfrac{i}{h}\tau\big), \qquad q_i' := \gamma\big(\tfrac{(i+s)}{h}\tau\big), \qquad \forall i = 0, ..., h-1,$$

and, for all $i = 0, ..., h-1$ and $t \in [0, \tau/h]$,

$$h_s(\gamma)\big(\tfrac{i}{h}\tau + t\big) = \begin{cases} \gamma_{q_i, q_i', s\tau/h}(t), & \text{if } t \in [0, s\tau/h], \\ \gamma(i\tau + t) & \text{if } t \in [s\tau/h, \tau/h]. \end{cases}$$

Here, we have adopted the notation of Section 3, denoting by $\gamma_{q_i, q_i', s\tau/h}$ the non-degenerate unique action minimizer given by Proposition 3.1. Notice that h_0 is the identity, and the image of h_1 is contained in the image of the embedding ι_h defined in (3.4). Moreover,

$$\tfrac{\mathrm{d}}{\mathrm{d}s} \mathcal{S}_k(h_s(\gamma)) \leq 0, \qquad \forall \gamma \in \mathcal{W},$$

that is, the homotopy h_s preserves the sublevels of the free-period action functional.

We choose h large enough so that $\zeta_1, ..., \zeta_r \in \mathcal{W}$. In particular, there exist points $(\boldsymbol{q}_i, \tau_i) \in \Delta_{h,\rho} \times (0, \epsilon)$ such that

$$\iota_h(\boldsymbol{q}_i, \tau_i) = \zeta_i, \qquad \forall i = 1, ..., r.$$

We set $c_i := \mathcal{S}_k(\zeta_i) = S_k(\boldsymbol{q}_i, \tau_i)$. Let us apply Lemma 4.3, which gives an integer $m_0 \in \mathbb{N}$ with the following property. For each $i \in \{1, ..., r\}$ and $m \geq m_0$, there exists a neighborhood $U_{i,m}$ of the critical circle of $(\boldsymbol{q}_i^m, \tau_i)$ such that the inclusion induces the injective map among connected components

$$\pi_0(\{S_k < mc_i\}) \hookrightarrow \pi_0(\{S_k < mc_i\} \cup U_{i,m}). \tag{4.5}$$

Now, by Lemma 4.5, $c(n,k) \to -\infty$ as $n \to \infty$. Therefore, if we choose $n \in \mathbb{N}$ large enough, we have that

$$\operatorname{crit}(\mathcal{S}_k) \cap \mathcal{S}_k^{-1}(c(n,k)) = \left\{\zeta_{i_1}^{m_1}, ..., \zeta_{i_u}^{m_u}\right\}$$

for some $i_1, ..., i_u \in \{1, ..., r\}$ and $m_1, ..., m_u \in \mathbb{N}$ such that

$$\min\{m_1, ..., m_u\} \geq m_0.$$

For each $v = 1, ..., u$, we fix a small enough open neighborhood $\mathcal{V}_v' \subset \mathcal{W}$ of the critical circle of $\zeta_{i_v}^{m_v}$ such that $h_1(\mathcal{V}_v') \subset \iota_{m_{i_v}h}(U_{i_v,m_{i_v}})$. We further choose a smaller open neighborhood $\mathcal{V}_v$ of the critical circle of $\zeta_{i_v}^{m_v}$ whose closure is contained in $\mathcal{V}_v'$. By Lemma 4.5, there exists a path $p \in \mathcal{P}(n,k)$ such that

$$p([0,1]) \subset \{S < c(n,k)\} \cup \mathcal{V}_1 \cup ... \cup \mathcal{V}_u.$$

Let $\chi : W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0,\infty) \to [0,1]$ be a continuous bump function that is identically equal to 1 on $\mathcal{V}_1 \cup ... \cup \mathcal{V}_u$ and is supported inside $\mathcal{V}_1' \cup ... \cup \mathcal{V}_u'$. We define the path $p' \in \mathcal{P}(n,k)$ by

$$p'(s) := h_{\chi(p(s))}(p(s)).$$

Notice that

$$p'([0,1]) \subset \{\mathcal{S}_k < c(n,k)\} \cup \iota_{m_1h}(U_{i_1,m_1}) \cup ... \cup \iota_{m_uh}(U_{i_u,m_u}).$$

This, together with equation (4.5), implies that there exists a path $p'' \in \mathcal{P}(n,k)$ such that $p''([0,1]) \subset \{\mathcal{S}_k < c(n,k)\}$, which contradicts the definition of the minimax value $c(n,k)$. This completes the proof. ∎

5. A Tonelli Lagrangian with few periodic orbits on low energy levels

If $L : \mathrm{T}M \to \mathbb{R}$ is a Tonelli Lagrangian with associated energy $E : \mathrm{T}M \to \mathbb{R}$, the Euler-Lagrange flow on any energy hypersurface $E^{-1}(k)$ is conjugate to the Hamiltonian flow of the dual Tonelli Hamiltonian $H : \mathrm{T}^*M \to \mathbb{R}$, which is given by (2.1), on the energy hypersurface $H^{-1}(k)$. We recall that the Hamiltonian vector field X_H generating the Hamiltonian flow is defined by $\mathrm{d}q \wedge \mathrm{d}p(X_H, \cdot\,) = \mathrm{d}H$. Moreover, $H(q,p) = E(q, H_p(q,p))$ for all $(q,p) \in \mathrm{T}^*M$, and therefore

$$\min E = \min H,$$
$$e_0(L) = e_0(H) := \min\left\{k \in \mathbb{R} \;\middle|\; \pi(H^{-1}(k)) = M\right\},$$

where $\pi : \mathrm{T}^*M \to M$ denotes the projection onto the base of the cotangent bundle.

We want to show that the assertions of Theorems 1.1 and 1.2 do not necessarily hold on the energy range $[\min E, e_0(L)]$. We will provide a counterexample for these statements in the Hamiltonian formulation. We fix two positive real numbers $r_1 < r_2$ such that their quotient r_1/r_2 is irrational, a real number $R > r_2$, and a smooth monotone function $\chi : [0,\infty) \to [0,\infty)$ such that $\chi(x) = x$ for all $x \in [0, r_2]$, and $\chi(x) = R$ for all $x \geq R$. We define a Tonelli Hamiltonian $H : \mathrm{T}^*\mathbb{R}^2 \to \mathbb{R}$ by

$$H(q_1, q_2, p_1, p_2) = \frac{1}{2}\left(\frac{\chi(|q_1|^2) + |p_1|^2}{r_1} + \frac{\chi(|q_2|^2) + |p_2|^2}{r_2}\right).$$

Notice that

$$0 = \min H < e_0(H) = \frac{1}{2}\left(\frac{R}{r_1} + \frac{R}{r_2}\right).$$

If we identify $\mathrm{T}^*\mathbb{R}^2$ with $\mathbb{C}^2$ by means of the map $(q_1, q_2, p_1, p_2) \mapsto (x_1+iy_1, x_2+iy_2)$, the Hamiltonian flow of H on the open polydisk $B(r_1) \times B(r_2) \subset \mathbb{C}^2$ is given by the unitary linear map

$$\phi_H^t = \begin{pmatrix} e^{-it/r_1} & 0 \\ 0 & e^{-it/r_2} \end{pmatrix}.$$

Since r_1/r_2 is irrational, for each $k \in (0, 1/2)$ there exists only two periodic orbits of the Hamiltonian flow ϕ_H^t with energy k. These orbits are given by

$$\Gamma_k : \mathbb{R}/2\pi r_1\mathbb{Z} \to \mathbb{C}^2, \qquad \Gamma_k(t) = \phi_H^t\big(\sqrt{2r_1k}, 0\big),$$
$$\Psi_k : \mathbb{R}/2\pi r_2\mathbb{Z} \to \mathbb{C}^2, \qquad \Psi_k(t) = \phi_H^t\big(0, \sqrt{2r_2k}\big).$$

Their Maslov index, which can be computed as in [Maz15, example 3.5], is given by

$$\mathrm{mas}(\Gamma_k) = 2\big(\lfloor r_1/r_2 \rfloor + 1\big),$$
$$\mathrm{mas}(\Psi_k) = 2\big(\lfloor r_2/r_1 \rfloor + 1\big).$$

Let $\pi : \mathrm{T}^*\mathbb{R}^2 \to \mathbb{R}$ be the base projection. The curves $\gamma_k := \pi \circ \Gamma_k$ and $\psi_k := \pi \circ \Psi_k$ are the only periodic orbits of the Lagrangian system of L with energy k. The Morse index Theorem for Tonelli Lagrangians (see, e.g., [Maz15, Theorem 4.1]) implies that the Morse index of the fixed-period Lagrangian action functional at a periodic orbit is equal to the Maslov index of its corresponding Hamiltonian periodic orbit. Since the Morse index of the free-period action functional at a periodic orbit is larger than or equal to the corresponding Morse index of the fixed-period action functional, and since $\mathrm{mas}(\Gamma_k) > 0$ and $\mathrm{mas}(\Psi_k) > 0$, we infer that γ_k and ψ_k are not local minima of the free-period action functional of L at energy k. Finally, notice that H does not depend on the base variables (q_1, q_2) in the region where $\min\{|q_1|^2, |q_2|^2\} > R$. Therefore, we can replace the configuration space with the 2-torus

$$\mathbb{T}^2 = ([-R, R]/\{-R, R\}) \times ([-R, R]/\{-R, R\}),$$

and see H as a Tonelli Hamiltonian of the form $H : \mathrm{T}^*\mathbb{T}^2 \to \mathbb{R}$. Equivalently we can see L as a Tonelli Lagrangian of the form $L : \mathrm{T}\mathbb{T}^2 \to \mathbb{R}$. Such Tonelli Lagrangian violates the assertions of Theorems 1.1 and 1.2 in the energy range $(0, 1/2) \subset [\min E, e_0(L)]$.

6. Tonelli setting with a general magnetic form

Let M be a closed manifold, $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian, and σ a closed 2-form on M. The pair (L, σ) defines a flow on the tangent bundle $\mathrm{T}M$, which we call the **Euler-Lagrange flow** of (L, σ), as follows: for each open subset $U \subset M$ such that $\sigma|_U$ admits a primitive θ, the Euler-Lagrange flow of (L, σ) coincides with the usual Euler-Lagrange flow of $L + \theta$ within $\mathrm{T}U$. This is a good definition, since the Euler-Lagrange flow of the Lagrangian L does not change if we add a closed 1-form to L. Since the energy function $E : \mathrm{T}M \to \mathbb{R}$ associated to L is also the energy function associated to $L + \theta$, the Euler-Lagrange flow of (L, σ) preserves E. This dynamics can be equivalently described in the Hamiltonian formulation as the Hamiltonian flow of the Tonelli Hamiltonian dual to L computed with respect to the twisted symplectic form $\mathrm{d}q \wedge \mathrm{d}p - \pi^*\sigma$ on T^*M, where $\pi : \mathrm{T}^*M \to M$ is the base projection.

The periodic orbits of the Euler-Lagrangian flow of (L,σ) on the energy hypersurface $E^{-1}(k)$ are in one-to-one correspondence with the zeroes of the action 1-form η_k on $W^{1,2}(\mathbb{R}/\mathbb{Z},M)\times(0,\infty)$, which is given by

$$\eta_k(\Gamma,\tau)\big[(\Psi,\mu)\big] = \mathrm{d}\mathcal{S}_k(\Gamma,\tau)\big[(\Psi,\mu)\big] + \int_0^1 \sigma_{\Gamma(t)}(\dot\Gamma(t),\Psi(t))\,\mathrm{d}t.$$

Here, we have denoted by $\mathcal{S}_k : W^{1,2}(\mathbb{R}/\mathbb{Z},M)\times(0,\infty)\to\mathbb{R}$ the free-period action functional of L at energy k. When the 2-form σ admits a primitive θ on M, the action 1-form η_k coincides with the differential of the free-period action functional with energy k of the Lagrangian $L+\theta$. We refer the reader to [AB14, AB15a, AB15b] and to the references therein for more background on the action 1-form. From now on, we assume that M is a closed surface, i.e.,

$$\dim(M) = 2.$$

Given a zero (Γ,τ) of the action 1-form, there exists a neighborhood $U\subset M$ of $\Gamma(\mathbb{R}/\mathbb{Z})$ such that $\sigma|_U$ is exact. In particular, on a neighborhood of (Γ,τ), η_k is the differential of the free-period action functional at energy k of the Lagrangian $L+\theta$, for any primitive θ of $\sigma|_U$.

We fix $k\in(e_0(L),c_0(L))$ and define the space $\mathcal{M}(n)$ as in Section 2, except that in condition (D1) we now require the restriction $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]}$ to be a unique local minimizer with energy k for the Lagrangian $L+\theta$, for some primitive θ of σ defined on a neighborhood of $\gamma_i|_{[\tau_{i,j},\tau_{i,j+1}]}$. Given a multicurve $\boldsymbol{\gamma}=(\gamma_1,...,\gamma_m)\in\mathcal{M}(n)$ we can find a sequence of embedded multicurves $\boldsymbol{\gamma}_\alpha\in\mathcal{M}(n)$ such that $\boldsymbol{\gamma}_\alpha\to\boldsymbol{\gamma}$ as $\alpha\to\infty$. By the definition of $\mathcal{M}(n)$, each multicurve $\boldsymbol{\gamma}_\alpha$ is the boundary of a possibly disconnected, embedded, oriented, compact surface $\Sigma_\alpha\subset M$ whose orientation agrees with the one of M. This allows to define a free-period action functional $\mathcal{S}_k:\mathcal{M}(n)\to\mathbb{R}$ by

$$\mathcal{S}_k(\boldsymbol{\gamma}) = \sum_{i=1}^m \left(\int_0^{\tau_i} L(\gamma_i(t),\dot\gamma_i(t))\,\mathrm{d}t + \tau_i k\right) + \int_\Sigma \sigma,$$

where

$$\Sigma = \bigcap_{\beta\in\mathbb{N}}\bigcup_{\alpha\geq\beta}\Sigma_\alpha.$$

The same arguments as in the proofs of Lemmas 2.8 and 2.9 imply that a multicurve $\boldsymbol{\gamma}\in\mathcal{M}(n-3)$ that is a global minimum of $\mathcal{S}_k$ over $\mathcal{M}(n)$ is embedded, and its connected components lift to periodic orbits of the Euler-Lagrange flow of (L,σ) on $E^{-1}(k)$.

For every finite covering space $M'\to M$, we can lift the Lagrangian systems of (L,σ) to a Lagrangian system (L',σ') on the configuration space M'. We denote by $\mathcal{M}'(n)$ and $\mathcal{S}'_k$ the space of multicurves and the free-period action functional associated to (L',σ'). We define $e(L,\sigma)$ as the supremum of $e(M',L,\sigma)\in\mathbb{R}$ over all the finite covering spaces $M'\to M$, where

$$e(M',L,\sigma) := \sup\Big\{k\geq e_0(L)\;\Big|\;\inf_{\mathcal{M}'(n)}\mathcal{S}'_k<0 \text{ for some } n\in\mathbb{N}\Big\}.$$

If σ is exact, $e(L,\sigma)$ coincides with the Mañé critical value of the abelian cover $c_0(L+\theta)$. The arguments in Sections 2.2 and 2.3 can be carried over for the Lagrangian system of (L,σ), and prove the following generalization of Theorem 1.1.

Theorem 6.1. *Let M be a closed surface, $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian, and σ a 2-form on M. For every $k \in (e_0(L), e(L,\sigma))$, the Lagrangian system of (L,σ) possesses a periodic orbit γ_k with energy k that is a local minimum of a local primitive of the action 1-form η_k. Moreover, γ_k lifts to a simple closed curve in some finite covering of M.* ■

Remark 6.2. It may happen that the interval $(e_0(L), e(L,\sigma))$ is empty, and in such case the assertion of Theorem 6.1 becomes void. However, when L has the form of a kinetic energy $L(q,v) = \frac{1}{2} g_q(v,v)$ for some Riemannian metric g, we have $e(L,\sigma) > e_0(L)$ if and only if σ is **oscillating**, that is, σ changes sign on M (see [AB15a, Lemma 6.4]).

If M is a closed surface other than the 2-sphere, the 2-form σ lifts to an exact form $\mathrm{d}\theta$ on the universal cover $\widetilde{M}$ of M. We define

$$e_*(L,\sigma) := \min\big\{e(L,\sigma), c(\widetilde{L}+\theta)\big\}.$$

Here, $\widetilde{L} : \mathrm{T}\widetilde{M} \to \mathbb{R}$ is the lift of L, and $c(\widetilde{L}+\theta)$ is the Mañé critical value of the Lagrangian $\widetilde{L}+\theta$. By combining the arguments of Sections 3 and 4.1 with the proofs of the main results in [AB14, AB15a, AB15b], we obtain the following generalization of Theorem 1.2.

Theorem 6.3. *Let M be a closed surface, $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian, and σ a 2-form on M.*

- *If $M \neq S^2$ or σ is exact, for almost every $k \in (e_0(L), e_*(L,\sigma))$ the Lagrangian system of (L,σ) possesses infinitely many periodic orbits with energy k.*
- *If $M = S^2$ and σ is not exact, for almost every $k \in (e_0(L), e(L,\sigma))$ the Lagrangian system of (L,σ) possesses two periodic orbits with energy k.*

■

After the first version of this paper appeared online, the authors, together with Abbondandolo, Benedetti, and Taimanov, completed the case of the 2-sphere in Theorem 6.3, showing that, when $M = S^2$, for almost every $k \in (e_0(L), e(L,\sigma))$ the Lagrangian system of (L,σ) possesses infinitely many periodic orbits with energy k, see [AAB$^+$16].

References

[AAB$^+$16] A. Abbondandolo, L. Asselle, G. Benedetti, M. Mazzucchelli, and I. A. Taimanov, *The multiplicity problem for periodic orbits of magnetic flows on the 2-sphere*, arXiv:1608.03144, 2016.

[AB14] L. Asselle and G. Benedetti, *The Lusternik-Fet theorem for autonomous Tonelli Hamiltonian systems on twisted cotangent bundles*, arXiv:1412.0531, to appear in J. Topol. Anal., 2014.

[AB15a] ———, *Infinitely many periodic orbits in non-exact oscillating magnetic fields on surfaces with genus at least two for almost every low energy level*, Calc. Var. Partial Differ. Equ. **54** (2015), no. 2, 1525–1545.

[AB15b] ———, *Periodic orbits in oscillating magnetic fields on T^2*, arXiv:1510.00152, 2015.

[Abb13] A. Abbondandolo, *Lectures on the free period Lagrangian action functional*, J. Fixed Point Theory Appl. **13** (2013), no. 2, 397–430.

[AMMP14] A. Abbondandolo, L. Macarini, M. Mazzucchelli, and G. P. Paternain, *Infinitely many periodic orbits of exact magnetic flows on surfaces for almost every subcritical energy level*, arXiv:1404.7641, to appear in J. Eur. Math. Soc. (JEMS), 2014.

[AMP15] A. Abbondandolo, L. Macarini, and G. P. Paternain, *On the existence of three closed magnetic geodesics for subcritical energies*, Comment. Math. Helv. **90** (2015), no. 1, 155–193.

[AP94] M. Abate and G. Patrizio, *Finsler metrics – a global approach*, Lecture Notes in Mathematics, vol. 1591, Springer-Verlag, Berlin, 1994.

[AS09] A. Abbondandolo and M. Schwarz, *A smooth pseudo-gradient for the Lagrangian action functional*, Adv. Nonlinear Stud. **9** (2009), no. 4, 597–623.

[Ban80] V. Bangert, *Closed geodesics on complete surfaces*, Math. Ann. **251** (1980), no. 1, 83–96.

[BCS00] D. Bao, S.-S. Chern, and Z. Shen, *An introduction to Riemann-Finsler geometry*, Graduate Texts in Mathematics, vol. 200, Springer-Verlag, New York, 2000.

[CDI97] G. Contreras, J. Delgado, and R. Iturriaga, *Lagrangian flows: the dynamics of globally minimizing orbits. II*, Bol. Soc. Brasil. Mat. (N.S.) **28** (1997), no. 2, 155–196.

[CI99] G. Contreras and R. Iturriaga, *Global minimizers of autonomous Lagrangians*, 22º Colóquio Brasileiro de Matemática, IMPA, Rio de Janeiro, 1999.

[CIPP98] G. Contreras, R. Iturriaga, G. P. Paternain, and M. Paternain, *Lagrangian graphs, minimizing measures and Mañé's critical values*, Geom. Funct. Anal. **8** (1998), no. 5, 788–809.

[CMP04] G. Contreras, L. Macarini, and G. P. Paternain, *Periodic orbits for exact magnetic flows on surfaces*, Int. Math. Res. Not. (2004), no. 8, 361–387.

[Con06] G. Contreras, *The Palais-Smale condition on contact type energy levels for convex Lagrangian systems*, Calc. Var. Partial Differ. Equ. **27** (2006), no. 3, 321–395.

[Fat97] A. Fathi, *Théorème KAM faible et théorie de Mather sur les systèmes lagrangiens*, C. R. Acad. Sci. Paris Sér. I Math. **324** (1997), no. 9, 1043–1046.

[Fat08] ———, *Weak KAM theorem in Lagrangian dynamics*, Cambridge Univ. Press, forthcoming, preliminary version number 10, 2008.

[GM69] D. Gromoll and W. Meyer, *Periodic geodesics on compact Riemannian manifolds*, J. Differ. Geom. **3** (1969), 493–510.

[Mañ97] R. Mañé, *Lagrangian flows: the dynamics of globally minimizing orbits*, Bol. Soc. Brasil. Mat. (N.S.) **28** (1997), no. 2, 141–153.

[Mat91a] J. N. Mather, *Action minimizing invariant measures for positive definite Lagrangian systems*, Math. Z. **207** (1991), no. 2, 169–207.

[Mat91b] ———, *Variational construction of orbits of twist diffeomorphisms*, J. Amer. Math. Soc. **4** (1991), no. 2, 207–263.

[Mat93] ———, *Variational construction of connecting orbits*, Ann. Inst. Fourier (Grenoble) **43** (1993), no. 5, 1349–1386.

[Maz11] M. Mazzucchelli, *The Lagrangian Conley conjecture*, Comment. Math. Helv. **86** (2011), no. 1, 189–246.

[Maz12] ———, *Critical point theory for Lagrangian systems*, Progress in Mathematics, vol. 293, Birkhäuser/Springer Basel AG, Basel, 2012.

[Maz14] ———, *On the multiplicity of isometry-invariant geodesics on product manifolds*, Algebr. Geom. Topol. **14** (2014), no. 1, 135–156.

[Maz15] ———, *The Morse index of Chaperon's generating families*, arXiv:1507.08344, to appear in Publ. Mat. Urug., 2015.

[Mil63] J. Milnor, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.

[MM15] L. Macarini and M. Mazzucchelli, *Isometry-invariant geodesics and the fundamental group, II*, arXiv:1504.05685, 2015.

[Sor15] A. Sorrentino, *Action-minimizing methods in Hamiltonian dynamics: an introduction to Aubry-Mather theory*, Mathematical Notes, vol. 50, Princeton University Press, Princeton, NJ, 2015.

[Tai91] I. A. Taimanov, *Non-self-itersecting closed extremals of multivalued or not everywhere positive functionals*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 2, 367–383.

[Tai92a] ———, *Closed extremals on two-dimensional manifolds*, Uspekhi Mat. Nauk **47** (1992), no. 2(284), 143–185, 223.

[Tai92b] ———, *Closed non-self-intersecting extremals of multivalued functionals*, Sibirsk. Mat. Zh. **33** (1992), no. 4, 155–162, 223.

[Ton34] L. Tonelli, *Su gli integrali del calcolo delle variazioni in forma ordinaria*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2) **3** (1934), no. 3-4, 401–450.

Ruhr Universität Bochum, Fakultät für Mathematik
Gebäude NA 4/33, D-44801 Bochum, Germany
E-mail address: luca.asselle@ruhr-uni-bochum.de

CNRS, École Normale Supérieure de Lyon, UMPA
69364 Lyon Cedex 07, France
E-mail address: marco.mazzucchelli@ens-lyon.fr